

Resource Summary Report

Generated by [RRID](#) on Aug 9, 2026

STS2 / TULA (internal) antibody

RRID:AB_10726216

Type: Antibody

Proper Citation

GeneTex Cat# GTX89140, RRID:AB_10726216

Antibody Information

URL: http://antibodyregistry.org/AB_10726216

Proper Citation: GeneTex Cat# GTX89140, RRID:AB_10726216

Target Antigen: STS2 / TULA (internal) antibody

Host Organism: goat

Clonality: polyclonal

Comments: Discontinued; manufacturer recommendations: Preliminary experiments gave an approx 48kDa band in lysates of B-cell line Daudi after 0.1µg/ml antibody staining. Please note that currently we cannot find an explanation in the literature for the band we observe given the calculated size of 74.1kDa according to NP_061834.1 and 69.8kDa according to NP_001001895.1. The 48kDa band was successfully blocked by incubation with the immunizing peptide. We would appreciate any feedback from people in the field - have any results been reported with other antibodies/lysates? Have any further splice variants/modified forms been reported? In paraffin embedded Human Spleen shows cytoplasm staining of T-cells eccentrically gathered around an artery. Recommended concentration, 2.5µg/ml. This antibody is expected to recognize both reported

Antibody Name: STS2 / TULA (internal) antibody

Description: This polyclonal targets STS2 / TULA (internal) antibody

Target Organism: canine, dog, human

Antibody ID: AB_10726216

Vendor: GeneTex

Catalog Number: GTX89140

Record Creation Time: 20250819T224027+0000

Ratings and Alerts

No rating or validation information has been found for STS2 / TULA (internal) antibody.

Warning: Discontinued at GeneTex

Discontinued; manufacturer recommendations: Preliminary experiments gave an approx 48kDa band in lysates of B-cell line Daudi after 0.1µg/ml antibody staining. Please note that currently we cannot find an explanation in the literature for the band we observe given the calculated size of 74.1kDa according to NP_061834.1 and 69.8kDa according to NP_001001895.1. The 48kDa band was successfully blocked by incubation with the immunizing peptide. We would appreciate any feedback from people in the field - have any results been reported with other antibodies/lysates? Have any further splice variants/modified forms been reported? In paraffin embedded Human Spleen shows cytoplasm staining of T-cells eccentrically gathered around an artery. Recommended concentration, 2.5µg/ml. This antibody is expected to recognize both reported

Data and Source Information

Source: [Antibody Registry](#)

Usage and Citation Metrics

We have not found any literature mentions for this resource.