

Resource Summary Report

Generated by [RRID](#) on Aug 8, 2026

Epstein-Barr virus Bcl-2 antibody [0235]

RRID:AB_11166795

Type: Antibody

Proper Citation

GeneTex Cat# GTX36645, RRID:AB_11166795

Antibody Information

URL: http://antibodyregistry.org/AB_11166795

Proper Citation: GeneTex Cat# GTX36645, RRID:AB_11166795

Target Antigen: Epstein-Barr virus Bcl-2 antibody [0235]

Host Organism: mouse

Clonality: monoclonal

Comments: Discontinued; manufacturer recommendations: Recommended Starting Dilutions: For ICC/IF: Use at 1:10-1:50; For WB: Use at 1:10-1:50; For IHC: Use at 1:10-1:50; For ELISA: Use at 1:20-1:200; Not yet tested in other applications. Optimal dilutions should be determined experimentally by the researcher.

For immunostaining cells containing EBNA-1 with clone 0211, slides must be fixed in methanol/acetone (3:7) at -20 °C. After drying at room temperature, slides should be stored at 4°C (1 month) or -20°C (6-12 months). A highly amplified assay must be used to detect the low amount of EBNA expressed, i.e. Monoclonal antibody + Goat Anti-Mouse IgG + Rabbit Anti-Goat IgG-FITC conj., Immunohistochemistry, Immunoprecipitation, Western Blot. The usefulness of this product in other applications has not been determined., IHC, IP, WB;

Antibody Name: Epstein-Barr virus Bcl-2 antibody [0235]

Description: This monoclonal targets Epstein-Barr virus Bcl-2 antibody [0235]

Target Organism: virus

Antibody ID: AB_11166795

Vendor: GeneTex

Catalog Number: GTX36645

Record Creation Time: 20250819T221452+0000

Record Last Update: 20250820T013503+0000

Ratings and Alerts

No rating or validation information has been found for Epstein-Barr virus Bcl-2 antibody [0235].

Warning: Discontinued at GeneTex

Discontinued; manufacturer recommendations: Recommended Starting Dilutions: For ICC/IF: Use at 1:10-1:50; For WB: Use at 1:10-1:50; For IHC: Use at 1:10-1:50; For ELISA: Use at 1:20-1:200; Not yet tested in other applications. Optimal dilutions should be determined experimentally by the researcher.

For immunostaining cells containing EBNA-1 with clone 0211, slides must be fixed in methanol/acetone (3:7) at -20 °C. After drying at room temperature, slides should be stored at 4°C (1 month) or -20°C (6-12 months). A highly amplified assay must be used to detect the low amount of EBNA expressed, i.e. Monoclonal antibody + Goat Anti-Mouse IgG + Rabbit Anti-Goat IgG-FITC conj., Immunohistochemistry, Immunoprecipitation, Western Blot. The usefulness of this product in other applications has not been determined., IHC, IP, WB;

Data and Source Information

Source: [Antibody Registry](#)

Usage and Citation Metrics

We have not found any literature mentions for this resource.