

Resource Summary Report

Generated by [RRID](#) on Sep 10, 2026

Blimp-1 Antibody

RRID:AB_11167295

Type: Antibody

Proper Citation

GeneTex Cat# GTX30605, RRID:AB_11167295

Antibody Information

URL: http://antibodyregistry.org/AB_11167295

Proper Citation: GeneTex Cat# GTX30605, RRID:AB_11167295

Target Antigen: Blimp-1

Host Organism: mouse

Clonality: monoclonal

Comments: Discontinued; manufacturer recommendations: IgG1; IgG1 Recognizes a band ~98 kD and may recognize one ~80 kDa (the beta form). Though this production of antibody has not been tested, previous lots have been used for immunohistochemistry, immunocytochemistry, and immunofluorescence on formalin-fixed paraffin-embedded or frozen sections. Antigen retrieval is recommended (EDTA buffer, microwave). This antibody demonstrates nuclear staining. This antibody does not immunoprecipitate. Suggested working dilutions: * Western Blot - 1:1,000-1:5,000 Immunocytochemistry - (+) previous supernatant tested at 1:50-1:100 Immunofluorescence - (+) previous supernatant tested at 1:200 with tyramide amplification Immunohistochemistry - (+) previous supernatant tested at 1:200 with tyramide amplification *The investigator should determine t

Antibody Name: Blimp-1 Antibody

Description: This monoclonal targets Blimp-1

Target Organism: mouse, human

Antibody ID: AB_11167295

Vendor: GeneTex

Catalog Number: GTX30605

Record Creation Time: 20231110T060328+0000

Record Last Update: 20260829T075939+0000

Ratings and Alerts

No rating or validation information has been found for Blimp-1 Antibody.

Warning: Discontinued at GeneTex

Discontinued; manufacturer recommendations: IgG1; IgG1 Recognizes a band ~98 kD and may recognize one ~80 kDa (the beta form). Though this production of antibody has not been tested, previous lots have been used for immunohistochemistry, immunocytochemistry, and immunofluorescence on formalin-fixed paraffin-embedded or frozen sections. Antigen retrieval is recommended (EDTA buffer, microwave). This antibody demonstrates nuclear staining. This antibody does not immunoprecipitate. Suggested working dilutions: * Western Blot - 1:1,000-1:5,000 Immunocytochemistry - (+) previous supernatant tested at 1:50-1:100 Immunofluorescence - (+) previous supernatant tested at 1:200 with tyramide amplification Immunohistochemistry - (+) previous supernatant tested at 1:200 with tyramide amplification *The investigator should determine t

Data and Source Information

Source: [Antibody Registry](#)

Usage and Citation Metrics

We have not found any literature mentions for this resource.