

Resource Summary Report

Generated by [RRID](#) on Aug 9, 2026

Prion protein PrP [7D9] antibody

RRID:AB_368110

Type: Antibody

Proper Citation

GeneTex Cat# GTX14219, RRID:AB_368110

Antibody Information

URL: http://antibodyregistry.org/AB_368110

Proper Citation: GeneTex Cat# GTX14219, RRID:AB_368110

Target Antigen: Prion protein PrP [7D9] antibody

Host Organism: mouse

Clonality: monoclonal

Comments: Discontinued; manufacturer recommendations: Immunohistochemistry - fixed; Western Blot; ELISA; Immunohistochemistry; ELISA, IHC (Formalin-fixed paraffin-embedded sections), Western blot. The usefulness of this product in other applications has not been determined., ELISA, IHC-P, WB, ELISA: Use at a dilution of 1/20,000 - 1/50,000. IHC-P: Use at a dilution of 1/100 - 1/500. Use at an assay dependent dilution. Retrieve antigens with 70 Formic acid for 10-30 minutes at room temperature or HIER (heat Induced Epitope Retrieval) before commencing with IHC staining protocol. WB: Use at a dilution of 1/10,000 - 1/25,000. Predicted molecular weight: 32 kDa. Not tested in other applications. Optimal dilutions/concentrations should be determined by the end user.

Antibody Name: Prion protein PrP [7D9] antibody

Description: This monoclonal targets Prion protein PrP [7D9] antibody

Target Organism: elk, hamster, cow, mouse, bovine, human, sheep

Antibody ID: AB_368110

Vendor: GeneTex

Catalog Number: GTX14219

Record Creation Time: 20250820T040854+0000

Record Last Update: 20250820T181036+0000

Ratings and Alerts

No rating or validation information has been found for Prion protein PrP [7D9] antibody.

Warning: Discontinued at GeneTex

Discontinued; manufacturer recommendations: Immunohistochemistry - fixed; Western Blot; ELISA; Immunohistochemistry; ELISA, IHC (Formalin-fixed paraffin-embedded sections), Western blot. The usefulness of this product in other applications has not been determined., ELISA, IHC-P, WB, ELISA: Use at a dilution of 1/20,000 - 1/50,000. IHC-P: Use at a dilution of 1/100 - 1/500. Use at an assay dependent dilution. Retrieve antigens with 70 Formic acid for 10-30 minutes at room temperature or HIER (heat Induced Epitope Retrieval) before commencing with IHC staining protocol. WB: Use at a dilution of 1/10,000 - 1/25,000. Predicted molecular weight: 32 kDa. Not tested in other applications. Optimal dilutions/concentrations should be determined by the end user.

Data and Source Information

Source: [Antibody Registry](#)

Usage and Citation Metrics

We have not found any literature mentions for this resource.