

Resource Summary Report

Generated by [RRID](#) on Sep 1, 2026

CD79a [HM47/A9] antibody

RRID:AB_368254

Type: Antibody

Proper Citation

GeneTex Cat# GTX20712, RRID:AB_368254

Antibody Information

URL: http://antibodyregistry.org/AB_368254

Proper Citation: GeneTex Cat# GTX20712, RRID:AB_368254

Target Antigen: CD79a [HM47/A9] antibody

Host Organism: mouse

Clonality: monoclonal

Comments: Discontinued; manufacturer recommendations: Immunohistochemistry, Immunoprecipitation, Western blot. The usefulness of this product in other applications has not been determined., IHC, IP, WB, This antibody may be diluted to a titer of 1:25-1:75 in an ABC method. However, depending on the staining conditions employed, we suggest that final dilution should be determined by the user. We suggest an incubation period of 30 minutes at room temperature. However, depending upon the fixation conditions and the staining system employed, optimal incubation should be determined by the user. Formalin fixed paraffin embedded tissue sections require high temperature antigen unmasking with 10 mM citrate buffer, pH 6.0 prior to immunostaining. IP: 1/50. WB: 1/50 - 1/100.; Immunohistochemistry; Immunoprecipitation; Western Blot

Antibody Name: CD79a [HM47/A9] antibody

Description: This monoclonal targets CD79a [HM47/A9] antibody

Target Organism: porcine, human

Antibody ID: AB_368254

Vendor: GeneTex

Catalog Number: GTX20712

Record Creation Time: 20250819T235137+0000

Ratings and Alerts

No rating or validation information has been found for CD79a [HM47/A9] antibody.

Warning: Discontinued at GeneTex

Discontinued; manufacturer recommendations: Immunohistochemistry, Immunoprecipitation, Western blot. The usefulness of this product in other applications has not been determined., IHC, IP, WB, This antibody may be diluted to a titer of 1:25-1:75 in an ABC method. However, depending on the staining conditions employed, we suggest that final dilution should be determined by the user. We suggest an incubation period of 30 minutes at room temperature. However, depending upon the fixation conditions and the staining system employed, optimal incubation should be determined by the user. Formalin fixed paraffin embedded tissue sections require high temperature antigen unmasking with 10 mM citrate buffer, pH 6.0 prior to immunostaining. IP: 1/50. WB: 1/50 - 1/100.; Immunohistochemistry; Immunoprecipitation; Western Blot

Data and Source Information

Source: [Antibody Registry](#)

Usage and Citation Metrics

We have not found any literature mentions for this resource.