

Resource Summary Report

Generated by [RRID](#) on Jul 30, 2026

[NGF beta antibody \[25623.1\]](#)

RRID:AB_368482

Type: Antibody

Proper Citation

GeneTex Cat# GTX10513, RRID:AB_368482

Antibody Information

URL: http://antibodyregistry.org/AB_368482

Proper Citation: GeneTex Cat# GTX10513, RRID:AB_368482

Target Antigen: NGF beta antibody [25623.1]

Host Organism: mouse

Clonality: monoclonal

Comments: Discontinued; manufacturer recommendations: IgG1; IgG1 Dot blot, ELISA, Inhibition assay. The usefulness of this product in other applications has not been determined., Dot, ELISA, Inhib, Dot: Use at a concentration of 1 - 3 µg/ml - determined using recombinant NGFB at 400 ng under non- and 200 ng under reducing conditions. ELISA: Use at a concentration of 1 - 2 µg/ml (100 µl/well) for capture ELISA, Monoclonal Anti-Human Nerve Growth Factor-Beta can be used as the capture antibody. Determined in combination with a biotinylated NGF-Beta polyclonal detection antibody at 50-100 ng/ml (100 µl/well), an ELISA for sample volumes of 100 µl with a range of 31.2 to 2000 pg/ml may be obtained. For capture ELISA, a working concentration of 1-2 µg/ml is determined to coat plates at 100 µl/well. Inhib: Use at a co

Antibody Name: NGF beta antibody [25623.1]

Description: This monoclonal targets NGF beta antibody [25623.1]

Target Organism: human

Antibody ID: AB_368482

Vendor: GeneTex

Catalog Number: GTX10513

Record Creation Time: 20250820T015455+0000

Ratings and Alerts

No rating or validation information has been found for NGF beta antibody [25623.1].

Warning: Discontinued at GeneTex

Discontinued; manufacturer recommendations: IgG1; IgG1 Dot blot, ELISA, Inhibition assay. The usefulness of this product in other applications has not been determined., Dot, ELISA, Inhib, Dot: Use at a concentration of 1 - 3 µg/ml - determined using recombinant NGFB at 400 ng under non- and 200 ng under reducing conditions. ELISA: Use at a concentration of 1 - 2 µg/ml (100 µl/well) for capture ELISA, Monoclonal Anti-Human Nerve Growth Factor-Beta can be used as the capture antibody. Determined in combination with a biotinylated NGF-Beta polyclonal detection antibody at 50-100 ng/ml (100 µl/well), an ELISA for sample volumes of 100 µl with a range of 31.2 to 2000 pg/ml may be obtained. For capture ELISA, a working concentration of 1-2 µg/ml is determined to coat plates at 100 µl/well. Inhib: Use at a co

Data and Source Information

Source: [Antibody Registry](#)

Usage and Citation Metrics

We have not found any literature mentions for this resource.