

Resource Summary Report

Generated by [RRID](#) on Aug 1, 2026

MUC1 [GP1.4] antibody (ready-to-use)

RRID:AB_368749

Type: Antibody

Proper Citation

GeneTex Cat# GTX20853, RRID:AB_368749

Antibody Information

URL: http://antibodyregistry.org/AB_368749

Proper Citation: GeneTex Cat# GTX20853, RRID:AB_368749

Target Antigen: MUC1 [GP1.4] antibody (ready-to-use)

Host Organism: mouse

Clonality: monoclonal

Comments: Discontinued; manufacturer recommendations: Immunohistochemistry - frozen; Immunohistochemistry - fixed; Immunohistochemistry; IHC (Frozen sections), IHC (Formalin-fixed paraffin-embedded sections). The usefulness of this product in other applications has not been determined., IHC-Fr, IHC-P, This antibody has been pretitered and quality controlled to work allon formalin-fixed paraffin-embedded as well as acetone fixed cryostat tissue sections. No further titration is required. We suggest an incubation period of 30-60 minutes at room temperature. However, depending upon the fixation conditions and the staining system employed, optimal incubation should be determined by the user. Enzymatic predigestion of paraffin embedded tissues is not necessary but in some tissues it may enhance the immunostaining.

Antibody Name: MUC1 [GP1.4] antibody (ready-to-use)

Description: This monoclonal targets MUC1 [GP1.4] antibody (ready-to-use)

Antibody ID: AB_368749

Vendor: GeneTex

Catalog Number: GTX20853

Record Creation Time: 20250820T010833+0000

Record Last Update: 20250820T100827+0000

Ratings and Alerts

No rating or validation information has been found for MUC1 [GP1.4] antibody (ready-to-use).

Warning: Discontinued at GeneTex

Discontinued; manufacturer recommendations: Immunohistochemistry - frozen; Immunohistochemistry - fixed; Immunohistochemistry; IHC (Frozen sections), IHC (Formalin-fixed paraffin-embedded sections). The usefulness of this product in other applications has not been determined., IHC-Fr, IHC-P, This antibody has been pretitered and quality controlled to work allon formalin-fixed paraffin-embedded as well as acetone fixed cryostat tissue sections. No further titration is required. We suggest an incubation period of 30-60 minutes at room temperature. However, depending upon the fixation conditions and the staining system employed, optimal incubation should be determined by the user. Enzymatic predigestion of paraffin embedded tissues is not necessary but in some tissues it may enhance the immunostaining.

Data and Source Information

Source: [Antibody Registry](#)

Usage and Citation Metrics

We have not found any literature mentions for this resource.