

Resource Summary Report

Generated by [RRID](#) on Sep 11, 2026

Cytokeratin 8 [C-51] antibody

RRID:AB_368920

Type: Antibody

Proper Citation

GeneTex Cat# GTX22531, RRID:AB_368920

Antibody Information

URL: http://antibodyregistry.org/AB_368920

Proper Citation: GeneTex Cat# GTX22531, RRID:AB_368920

Target Antigen: Cytokeratin 8 [C-51] antibody

Host Organism: mouse

Clonality: monoclonal

Comments: Discontinued; manufacturer recommendations: Western Blot; Immunohistochemistry - fixed; Immunoprecipitation; Immunohistochemistry; Immunocytochemistry; ICC, IHC-P, IP, WB, Immunocytochemistry, IHC (Formalin-fixed paraffin-embedded sections), Immunoprecipitation, Western blot. The usefulness of this product in other applications has not been determined., For ICC: Use at an assay dependant dilution. For IHC-P: Use at an assay dependant dilution. Perform enzymatic antigen retrieval before commencing with IHC staining protocol. For IP: Use at an assay dependant dilution. For WB: Use at an assay dependant dilution. Predicted molecular weight: 52.5 kDa. Not tested in other applications. Optimal dilutions/concentrations should be determined by the researcher.

Antibody Name: Cytokeratin 8 [C-51] antibody

Description: This monoclonal targets Cytokeratin 8 [C-51] antibody

Target Organism: donkey, porcine, cow, mouse, rabbit, bovine, human

Antibody ID: AB_368920

Vendor: GeneTex

Catalog Number: GTX22531

Record Creation Time: 20231110T081239+0000

Ratings and Alerts

No rating or validation information has been found for Cytokeratin 8 [C-51] antibody.

Warning: Discontinued at GeneTex

Discontinued; manufacturer recommendations: Western Blot; Immunohistochemistry - fixed; Immunoprecipitation; Immunohistochemistry; Immunocytochemistry; ICC, IHC-P, IP, WB, Immunocytochemistry, IHC (Formalin-fixed paraffin-embedded sections), Immunoprecipitation, Western blot. The usefulness of this product in other applications has not been determined., For ICC: Use at an assay dependant dilution. For IHC-P: Use at an assay dependant dilution. Perform enzymatic antigen retrieval before commencing with IHC staining protocol. For IP: Use at an assay dependant dilution. For WB: Use at an assay dependant dilution. Predicted molecular weight: 52.5 kDa. Not tested in other applications. Optimal dilutions/concentrations should be determined by the researcher.

Data and Source Information

Source: [Antibody Registry](#)

Usage and Citation Metrics

We have not found any literature mentions for this resource.