

Resource Summary Report

Generated by [RRID](#) on Aug 12, 2026

Caspase 8 [8CSP03] antibody

RRID:AB_368973

Type: Antibody

Proper Citation

GeneTex Cat# GTX23249, RRID:AB_368973

Antibody Information

URL: http://antibodyregistry.org/AB_368973

Proper Citation: GeneTex Cat# GTX23249, RRID:AB_368973

Target Antigen: Caspase 8 [8CSP03] antibody

Host Organism: mouse

Clonality: monoclonal

Comments: Discontinued; manufacturer recommendations: Western Blot; Immunohistochemistry - fixed; Immunoprecipitation; Immunohistochemistry; For IP (denatured use protein G): Use at a concentration of 2µg / mg of lysate. For WB: Use at a concentration of 2 - 4 µg / ml for 2hrs at room temperature. For IHC-P: Use at a concentration of 1 - 2 µg / ml for 30 min at room temperature. Staining of formalin-fixed tissues requires boiling tissue sections in 10mM citrate buffer, pH 6.0, for 10-20 min followed by cooling at RT for 20 min. Not tested in other applications. Optimal dilutions/concentrations should be determined by the researcher., IHC (Formalin-fixed paraffin-embedded sections), Immunoprecipitation, Western blot. The usefulness of this product in other applications has not been determined., IHC-P, IP, WB

Antibody Name: Caspase 8 [8CSP03] antibody

Description: This monoclonal targets Caspase 8 [8CSP03] antibody

Antibody ID: AB_368973

Vendor: GeneTex

Catalog Number: GTX23249

Record Creation Time: 20250820T070303+0000

Record Last Update: 20250821T013002+0000

Ratings and Alerts

No rating or validation information has been found for Caspase 8 [8CSP03] antibody.

Warning: Discontinued at GeneTex

Discontinued; manufacturer recommendations: Western Blot; Immunohistochemistry - fixed; Immunoprecipitation; Immunohistochemistry; For IP (denatured use protein G): Use at a concentration of 2µg / mg of lysate. For WB: Use at a concentration of 2 - 4 µg / ml for 2hrs at room temperature. For IHC-P: Use at a concentration of 1 - 2 µg / ml for 30 min at room temperature. Staining of formalin-fixed tissues requires boiling tissue sections in 10mM citrate buffer, pH 6.0, for 10-20 min followed by cooling at RT for 20 min. Not tested in other applications. Optimal dilutions/concentrations should be determined by the researcher., IHC (Formalin-fixed paraffin-embedded sections), Immunoprecipitation, Western blot. The usefulness of this product in other applications has not been determined., IHC-P, IP, WB

Data and Source Information

Source: [Antibody Registry](#)

Usage and Citation Metrics

We have not found any literature mentions for this resource.