

Resource Summary Report

Generated by [RRID](#) on Jul 29, 2026

Myogenin [SPM144] antibody

RRID:AB_368996

Type: Antibody

Proper Citation

GeneTex Cat# GTX15232, RRID:AB_368996

Antibody Information

URL: http://antibodyregistry.org/AB_368996

Proper Citation: GeneTex Cat# GTX15232, RRID:AB_368996

Target Antigen: Myogenin [SPM144] antibody

Host Organism: mouse

Clonality: monoclonal

Comments: Discontinued; manufacturer recommendations: Immunohistochemistry; Immunoprecipitation; Immunohistochemistry - fixed; Western Blot; IHC-P: Use at a dilution of 1/40 for 30 minutes at RT. Staining of formalin-fixed tissues requires boiling tissue sections in 10mM citrate buffer, pH 6.0 for 10 minutes followed by cooling at RT for 20 minutes. IP: Use at an assay dependent dilution. WB: Use at an assay dependent dilution. Predicted molecular weight: 25 kDa. Not tested in other applications. Optimal dilutions/concentrations should be determined by the end user., IHC (Formalin-fixed paraffin-embedded sections), Immunoprecipitation, Western blot. The usefulness of this product in other applications has not been determined., IHC-P, IP, WB

Antibody Name: Myogenin [SPM144] antibody

Description: This monoclonal targets Myogenin [SPM144] antibody

Target Organism: rat, mouse, human

Antibody ID: AB_368996

Vendor: GeneTex

Catalog Number: GTX15232

Record Creation Time: 20250819T222951+0000

Record Last Update: 20250820T022245+0000

Ratings and Alerts

No rating or validation information has been found for Myogenin [SPM144] antibody.

Warning: Discontinued at GeneTex

Discontinued; manufacturer recommendations: Immunohistochemistry; Immunoprecipitation; Immunohistochemistry - fixed; Western Blot; IHC-P: Use at a dilution of 1/40 for 30 minutes at RT. Staining of formalin-fixed tissues requires boiling tissue sections in 10mM citrate buffer, pH 6.0 for 10 minutes followed by cooling at RT for 20 minutes. IP: Use at an assay dependent dilution. WB: Use at an assay dependent dilution. Predicted molecular weight: 25 kDa. Not tested in other applications. Optimal dilutions/concentrations should be determined by the end user., IHC (Formalin-fixed paraffin-embedded sections), Immunoprecipitation, Western blot. The usefulness of this product in other applications has not been determined., IHC-P, IP, WB

Data and Source Information

Source: [Antibody Registry](#)

Usage and Citation Metrics

We have not found any literature mentions for this resource.