

Resource Summary Report

Generated by [RRID](#) on Jul 29, 2026

CDK2 [2B6] antibody

RRID:AB_369268

Type: Antibody

Proper Citation

GeneTex Cat# GTX22363, RRID:AB_369268

Antibody Information

URL: http://antibodyregistry.org/AB_369268

Proper Citation: GeneTex Cat# GTX22363, RRID:AB_369268

Target Antigen: CDK2 [2B6] antibody

Host Organism: mouse

Clonality: monoclonal

Comments: Discontinued; manufacturer recommendations: IHC: This antibody may be diluted to a titer of 1:25-1:50 in an ABC method. We suggest an incubation period of 30 minutes at room temperature. However, depending upon the fixation conditions and the staining system employed, optimal incubation conditions and antibody dilutions should be determined by the user. Formalin fixed paraffin embedded tissue sections require high temperature antigen unmasking with 1 mM EDTA buffer, pH 8.0 prior to immunostaining., IHC (Frozen sections), IHC (Formalin-fixed paraffin-embedded sections), Western blot. The usefulness of this product in other applications has not been determined., IHC-Fr, IHC-P, WB; Immunohistochemistry - frozen; Immunohistochemistry - fixed; Immunohistochemistry; Western Blot

Antibody Name: CDK2 [2B6] antibody

Description: This monoclonal targets CDK2 [2B6] antibody

Target Organism: rat, mouse, human

Antibody ID: AB_369268

Vendor: GeneTex

Catalog Number: GTX22363

Record Creation Time: 20250819T231701+0000

Ratings and Alerts

No rating or validation information has been found for CDK2 [2B6] antibody.

Warning: Discontinued at GeneTex

Discontinued; manufacturer recommendations: IHC: This antibody may be diluted to a titer of 1:25-1:50 in an ABC method. We suggest an incubation period of 30 minutes at room temperature. However, depending upon the fixation conditions and the staining system employed, optimal incubation conditions and antibody dilutions should be determined by the user. Formalin fixed paraffin embedded tissue sections require high temperature antigen unmasking with 1 mM EDTA buffer, pH 8.0 prior to immunostaining., IHC (Frozen sections), IHC (Formalin-fixed paraffin-embedded sections), Western blot. The usefulness of this product in other applications has not been determined., IHC-Fr, IHC-P, WB; Immunohistochemistry - frozen; Immunohistochemistry - fixed; Immunohistochemistry; Western Blot

Data and Source Information

Source: [Antibody Registry](#)

Usage and Citation Metrics

We found 1 mentions in open access literature.

Listed below are recent publications. The full list is available at [RRID](#) .

G??ttle P, et al. (2015) Oligodendroglial maturation is dependent on intracellular protein shuttling. The Journal of neuroscience : the official journal of the Society for Neuroscience, 35(3), 906.