

Resource Summary Report

Generated by [RRID](#) on Sep 11, 2026

CG antibody (ready-to-use)

RRID:AB_369401

Type: Antibody

Proper Citation

GeneTex Cat# GTX20932, RRID:AB_369401

Antibody Information

URL: http://antibodyregistry.org/AB_369401

Proper Citation: GeneTex Cat# GTX20932, RRID:AB_369401

Target Antigen: CG antibody (ready-to-use)

Host Organism: rabbit

Clonality: polyclonal

Comments: Discontinued; manufacturer recommendations: IgG; IgG Immunohistochemistry; Immunohistochemistry - frozen; Immunohistochemistry - fixed; This antibody has been pretitered and quality controlled to work allon formalin-fixed paraffin-embedded as well as acetone fixed cryostat tissue sections. No further titration is required. We suggest an incubation period of 30-60 minutes at room temperature. However, depending upon the fixation conditions and the staining system employed, optimal incubation conditions should be determined by the user. Although not required, proteolytic treatment may enhance the immunostaining in overfixed tissues. IHC (Frozen sections), IHC (Formalin-fixed paraffin-embedded sections).

Antibody Name: CG antibody (ready-to-use)

Description: This polyclonal targets CG antibody (ready-to-use)

Target Organism: human

Antibody ID: AB_369401

Vendor: GeneTex

Catalog Number: GTX20932

Record Creation Time: 20231110T081257+0000

Record Last Update: 20260901T043617+0000

Ratings and Alerts

No rating or validation information has been found for CG antibody (ready-to-use).

Warning: Discontinued at GeneTex

Discontinued; manufacturer recommendations: IgG; IgG Immunohistochemistry; Immunohistochemistry - frozen; Immunohistochemistry - fixed; This antibody has been pretitered and quality controlled to work allon formalin-fixed paraffin-embedded as well as acetone fixed cryostat tissue sections. No further titration is required. We suggest an incubation period of 30-60 minutes at room temperature. However, depending upon the fixation conditions and the staining system employed, optimal incubation conditions should be determined by the user. Although not required, proteolytic treatment may enhance the immunostaining in overfixed tissues. IHC (Frozen sections), IHC (Formalin-fixed paraffin-embedded sections).

Data and Source Information

Source: [Antibody Registry](#)

Usage and Citation Metrics

We have not found any literature mentions for this resource.