

Resource Summary Report

Generated by [RRID](#) on Aug 5, 2026

Topoisomerase II alpha antibody

RRID:AB_370102

Type: Antibody

Proper Citation

GeneTex Cat# GTX22987, RRID:AB_370102

Antibody Information

URL: http://antibodyregistry.org/AB_370102

Proper Citation: GeneTex Cat# GTX22987, RRID:AB_370102

Target Antigen: Topoisomerase II alpha antibody

Host Organism: rabbit

Clonality: polyclonal

Comments: Discontinued; manufacturer recommendations: IgG; IgG Immunofluorescence / Immunohistochemistry (formalin fixed paraffin embedded): assay dependent. Immunoprecipitation (with protein A, native and denatured): use at a concentration of 10µg / mg of lysate. Western Blotting: use at a concentration of 5 - 10µg / ml for 2hrs at RT. Not tested in other applications. Optimal dilutions / concentrations should be determined by the end user. By Western blot, this antibody detects a band of ~ 170 kDa, which corresponds to the predicted molecular weight of Topoisomerase II alpha. Immunofluorescence, IHC (Formalin-fixed paraffin-embedded sections), Immunoprecipitation, Western blot.; Immunoprecipitation; Immunofluorescence; Immunohistochemistry; Western Blot

Antibody Name: Topoisomerase II alpha antibody

Description: This polyclonal targets Topoisomerase II alpha antibody

Target Organism: human

Antibody ID: AB_370102

Vendor: GeneTex

Catalog Number: GTX22987

Record Creation Time: 20250820T043838+0000

Ratings and Alerts

No rating or validation information has been found for Topoisomerase II alpha antibody.

Warning: Discontinued at GeneTex

Discontinued; manufacturer recommendations: IgG; IgG Immunofluorescence / Immunohistochemistry (formalin fixed paraffin embedded): assay dependent. Immunoprecipitation (with protein A, native and denatured): use at a concentration of 10µg / mg of lysate. Western Blotting: use at a concentration of 5 - 10µg / ml for 2hrs at RT. Not tested in other applications. Optimal dilutions / concentrations should be determined by the end user. By Western blot, this antibody detects a band of ~ 170 kDa, which corresponds to the predicted molecular weight of Topoisomerase II alpha. Immunofluorescence, IHC (Formalin-fixed paraffin-embedded sections), Immunoprecipitation, Western blot.; Immunoprecipitation; Immunofluorescence; Immunohistochemistry; Western Blot

Data and Source Information

Source: [Antibody Registry](#)

Usage and Citation Metrics

We have not found any literature mentions for this resource.