

Resource Summary Report

Generated by [RRID](#) on Aug 4, 2026

MMP26 antibody

RRID:AB_370275

Type: Antibody

Proper Citation

GeneTex Cat# GTX11685, RRID:AB_370275

Antibody Information

URL: http://antibodyregistry.org/AB_370275

Proper Citation: GeneTex Cat# GTX11685, RRID:AB_370275

Target Antigen: MMP26 antibody

Host Organism: rabbit

Clonality: polyclonal

Comments: Discontinued; manufacturer recommendations: IgG; IgG Western Blot; WB: Use at a dilution of 1/1000. A starting dilution of 1/5000 is recommended for chemiluminescent substrates. By immunoblotting against the reduced protein, the antibody identifies bands at 30 kDa and 18 kDa (zymogen and active MMP26). MMP26 (matrilysin 2) does not appear to be produced by most normal quiescent cells, but the treatment of many cell types with the phorbol ester TPA or IL1 stimulates production of MMP26. Because of the low protein levels produced (pg/ml) concentration of cell culture media is often required to visualize the bands by immunoblotting. Note: Higher antibody dilutions may be necessary for non-human samples. Not tested in other applications. Optimal dilutions/concentrations should be determined by the end user. Western Blot.

Antibody Name: MMP26 antibody

Description: This polyclonal targets MMP26 antibody

Target Organism: human

Antibody ID: AB_370275

Vendor: GeneTex

Catalog Number: GTX11685

Record Creation Time: 20250820T001217+0000

Ratings and Alerts

No rating or validation information has been found for MMP26 antibody.

Warning: Discontinued at GeneTex

Discontinued; manufacturer recommendations: IgG; IgG Western Blot; WB: Use at a dilution of 1/1000. A starting dilution of 1/5000 is recommended for chemiluminescent substrates. By immunoblotting against the reduced protein, the antibody identifies bands at 30 kDa and 18 kDa (zymogen and active MMP26). MMP26 (matrilysin 2) does not appear to be produced by most normal quiescent cells, but the treatment of many cell types with the phorbol ester TPA or IL1 stimulates production of MMP26. Because of the low protein levels produced (pg/ml) concentration of cell culture media is often required to visualize the bands by immunoblotting. Note: Higher antibody dilutions may be necessary for non-human samples. Not tested in other applications. Optimal dilutions/concentrations should be determined by the end user. Western Blot.

Data and Source Information

Source: [Antibody Registry](#)

Usage and Citation Metrics

We have not found any literature mentions for this resource.