

Resource Summary Report

Generated by [RRID](#) on Aug 4, 2026

IL1-R2 [1.384.92.17.19] antibody

RRID:AB_371855

Type: Antibody

Proper Citation

GeneTex Cat# GTX13776, RRID:AB_371855

Antibody Information

URL: http://antibodyregistry.org/AB_371855

Proper Citation: GeneTex Cat# GTX13776, RRID:AB_371855

Target Antigen: IL1-R2 [1.384.92.17.19] antibody

Host Organism: mouse

Clonality: monoclonal

Comments: Discontinued; manufacturer recommendations: ELISA: Use at an assay dependent dilution. Can be used to measure natural human IL 1 Receptor antagonist levels in the sinovial fluid and in the serum as well as in cultured monocytes derived from patients with rheumatoid arthritis, in the range of 20 to 2000 pg/ml. See Towbin H et al. IA (Chemiluminescent): Use at an assay dependent dilution. IHC-Fr: Use at an assay dependent dilution. Tested on cryopreserved sections of rectal mucosal tissue from patients with shigellosis, and on frozen sections of tissue from children with tonsillitis. See references Raqib R et al. and Andersson J et al. Not tested in other applications. Optimal dilutions/concentrations should be determined by the end user., ELISA, Immunoassay, IHC (Frozen sections). The usefulness of this product in other applications

Antibody Name: IL1-R2 [1.384.92.17.19] antibody

Description: This monoclonal targets IL1-R2 [1.384.92.17.19] antibody

Antibody ID: AB_371855

Vendor: GeneTex

Catalog Number: GTX13776

Record Creation Time: 20250820T011506+0000

Record Last Update: 20250820T102548+0000

Ratings and Alerts

No rating or validation information has been found for IL1-R2 [1.384.92.17.19] antibody.

Warning: Discontinued at GeneTex

Discontinued; manufacturer recommendations: ELISA: Use at an assay dependent dilution. Can be used to measure natural human IL 1 Receptor antagonist levels in the synovial fluid and in the serum as well as in cultured monocytes derived from patients with rheumatoid arthritis, in the range of 20 to 2000 pg/ml. See Towbin H et al. IA (Chemiluminescent): Use at an assay dependent dilution. IHC-Fr: Use at an assay dependent dilution. Tested on cryopreserved sections of rectal mucosal tissue from patients with shigellosis, and on frozen sections of tissue from children with tonsillitis. See references Raqib R et al. and Andersson J et al. Not tested in other applications. Optimal dilutions/concentrations should be determined by the end user., ELISA, Immunoassay, IHC (Frozen sections). The usefulness of this product in other applications

Data and Source Information

Source: [Antibody Registry](#)

Usage and Citation Metrics

We have not found any literature mentions for this resource.