

Resource Summary Report

Generated by [RRID](#) on Aug 4, 2026

[IL7 antibody \[7417.111\]](#)

RRID:AB_371948

Type: Antibody

Proper Citation

GeneTex Cat# GTX10765, RRID:AB_371948

Antibody Information

URL: http://antibodyregistry.org/AB_371948

Proper Citation: GeneTex Cat# GTX10765, RRID:AB_371948

Target Antigen: IL7 antibody [7417.111]

Host Organism: mouse

Clonality: monoclonal

Comments: Discontinued; manufacturer recommendations: IgG1; IgG1 Western Blot; Block/Neutralize/Inhibit; Immunohistochemistry; ELISA; ELISA: Use at a concentration of 1 - 2 µg/ml. Neut: Use at an assay dependent dilution. WB: Use at a concentration of 1 - 2 µg/ml. Predicted molecular weight: 17 kDa. The exact concentration of antibody required to neutralize recombinant human IL7 activity is dependent on the cytokine concentration, cell type, growth conditions and the type of activity. The detection limit in immunoblotting for recombinant human IL7 is approximately 25 ng/lane and 300 ng/lane under nonreducing and reducing conditions, respectively. A working concentration of the capture antibody at 1-2 µg/ml (100µl/well) in combination with a biotinylated polyclonal detection antibody at 25-50 ng/ml (100µl/well) is recommend

Antibody Name: IL7 antibody [7417.111]

Description: This monoclonal targets IL7 antibody [7417.111]

Antibody ID: AB_371948

Vendor: GeneTex

Catalog Number: GTX10765

Record Creation Time: 20250819T230207+0000

Record Last Update: 20250820T040525+0000

Ratings and Alerts

No rating or validation information has been found for IL7 antibody [7417.111].

Warning: Discontinued at GeneTex

Discontinued; manufacturer recommendations: IgG1; IgG1 Western Blot; Block/Neutralize/Inhibit; Immunohistochemistry; ELISA; ELISA: Use at a concentration of 1 - 2 µg/ml. Neut: Use at an assay dependent dilution. WB: Use at a concentration of 1 - 2 µg/ml. Predicted molecular weight: 17 kDa. The exact concentration of antibody required to neutralize recombinant human IL7 activity is dependent on the cytokine concentration, cell type, growth conditions and the type of activity. The detection limit in immunoblotting for recombinant human IL7 is approximately 25 ng/lane and 300 ng/lane under nonreducing and reducing conditions, respectively. A working concentration of the capture antibody at 1-2 µg/ml (100µl/well) in combination with a biotinylated polyclonal detection antibody at 25-50 ng/ml (100µl/well) is recommend

Data and Source Information

Source: [Antibody Registry](#)

Usage and Citation Metrics

We have not found any literature mentions for this resource.