

# Resource Summary Report

Generated by [RRID](#) on Jul 30, 2026

## Integrin beta 5 antibody

RRID:AB\_371972

Type: Antibody

### Proper Citation

GeneTex Cat# GTX15459, RRID:AB\_371972

### Antibody Information

**URL:** [http://antibodyregistry.org/AB\\_371972](http://antibodyregistry.org/AB_371972)

**Proper Citation:** GeneTex Cat# GTX15459, RRID:AB\_371972

**Target Antigen:** Integrin beta 5 antibody

**Host Organism:** rabbit

**Clonality:** polyclonal

**Comments:** Discontinued; manufacturer recommendations: IgG; IgG Immunohistochemistry - fixed; Immunofluorescence; Immunohistochemistry; Immunoprecipitation; Western Blot; IF: Use at an assay dependent dilution. IHC-P: Use at a dilution of 1/50. Staining of formalin-fixed tissues requires boiling tissue sections in 10 mM citrate, pH 6.0 for 10 min followed by cooling at RT for 20 min. IP: Use at an assay dependent dilution. WB: Use at an assay dependent dilution. Predicted molecular weight: 95.9 kDa. Not tested in other applications. Optimal dilutions/concentrations should be determined by the end user. Immunofluorescence, IHC (Formalin-fixed paraffin-embedded sections), Immunoprecipitation, Western blot.

**Antibody Name:** Integrin beta 5 antibody

**Description:** This polyclonal targets Integrin beta 5 antibody

**Target Organism:** rat, hamster, mouse, human

**Antibody ID:** AB\_371972

**Vendor:** GeneTex

**Catalog Number:** GTX15459

**Record Creation Time:** 20250820T001016+0000

**Record Last Update:** 20250820T073334+0000

---

## Ratings and Alerts

No rating or validation information has been found for Integrin beta 5 antibody.

**Warning:** Discontinued at GeneTex

Discontinued; manufacturer recommendations: IgG; IgG Immunohistochemistry - fixed; Immunofluorescence; Immunohistochemistry; Immunoprecipitation; Western Blot; IF: Use at an assay dependent dilution. IHC-P: Use at a dilution of 1/50. Staining of formalin-fixed tissues requires boiling tissue sections in 10 mM citrate, pH 6.0 for 10 min followed by cooling at RT for 20 min. IP: Use at an assay dependent dilution. WB: Use at an assay dependent dilution. Predicted molecular weight: 95.9 kDa. Not tested in other applications. Optimal dilutions/concentrations should be determined by the end user. Immunofluorescence, IHC (Formalin-fixed paraffin-embedded sections), Immunoprecipitation, Western blot.

---

## Data and Source Information

**Source:** [Antibody Registry](#)

---

## Usage and Citation Metrics

We have not found any literature mentions for this resource.