

Resource Summary Report

Generated by [RRID](#) on Aug 4, 2026

Involucrin [0.N.341] antibody

RRID:AB_371977

Type: Antibody

Proper Citation

GeneTex Cat# GTX14505, RRID:AB_371977

Antibody Information

URL: http://antibodyregistry.org/AB_371977

Proper Citation: GeneTex Cat# GTX14505, RRID:AB_371977

Target Antigen: Involucrin [0.N.341] antibody

Host Organism: mouse

Clonality: monoclonal

Comments: Discontinued; manufacturer recommendations: Immunoprecipitation; ELISA; Western Blot; Immunofluorescence; Immunohistochemistry; Immunohistochemistry - fixed; ELISA, Immunofluorescence, IHC (Formalin-fixed paraffin-embedded sections), Immunoprecipitation, Western blot. The usefulness of this product in other applications has not been determined., ELISA, IF, IHC-P, IP, WB, ELISA: Use at an assay dependent dilution. IF: Use at an assay dependent dilution. IHC-P: Use at a concentration of 1 - 2 µg/ml. Perform enzymatic antigen retrieval before commencing with IHC staining protocol. For staining formalin-fixed tissues, digest sections with trypsin at 1mg/ml PBS, for 10 min at 37°C. Note: Enzyme digestion is better than citrate buffer based epitope unmasking. IP: Use 2 µg/mg protein lysate (Denatured verified; Use Protein

Antibody Name: Involucrin [0.N.341] antibody

Description: This monoclonal targets Involucrin [0.N.341] antibody

Target Organism: gorilla, monkey, rat, porcine, canine, chickenbird, human, dog

Antibody ID: AB_371977

Vendor: GeneTex

Catalog Number: GTX14505

Record Creation Time: 20250820T043632+0000

Ratings and Alerts

No rating or validation information has been found for Involucrin [O.N.341] antibody.

Warning: Discontinued at GeneTex

Discontinued; manufacturer recommendations: Immunoprecipitation; ELISA; Western Blot; Immunofluorescence; Immunohistochemistry; Immunohistochemistry - fixed; ELISA, Immunofluorescence, IHC (Formalin-fixed paraffin-embedded sections), Immunoprecipitation, Western blot. The usefulness of this product in other applications has not been determined., ELISA, IF, IHC-P, IP, WB, ELISA: Use at an assay dependent dilution. IF: Use at an assay dependent dilution. IHC-P: Use at a concentration of 1 - 2 µg/ml. Perform enzymatic antigen retrieval before commencing with IHC staining protocol. For staining formalin-fixed tissues, digest sections with trypsin at 1mg/ml PBS, for 10 min at 37°C. Note: Enzyme digestion is better than citrate buffer based epitope unmasking. IP: Use 2 µg/mg protein lysate (Denatured verified; Use Protein

Data and Source Information

Source: [Antibody Registry](#)

Usage and Citation Metrics

We have not found any literature mentions for this resource.