

Resource Summary Report

Generated by [RRID](#) on Sep 11, 2026

NKX3A antibody

RRID:AB_372507

Type: Antibody

Proper Citation

GeneTex Cat# GTX12044, RRID:AB_372507

Antibody Information

URL: http://antibodyregistry.org/AB_372507

Proper Citation: GeneTex Cat# GTX12044, RRID:AB_372507

Target Antigen: NKX3A antibody

Host Organism: rabbit

Clonality: polyclonal

Comments: Discontinued; manufacturer recommendations: IgG; IgG Immunohistochemistry; Other; Western Blot; IHC: Use at a dilution of 1/200 - 1/1000. WB: Use at a dilution of 1/200 - 1/1000. Detects a band of approximately 21.6 - 26.3 kDa. Preparation Instructions Method 1 for immunostaining and immunoblotting (Western blot): Remove a fixed amount of suspension and dissolve 1/5 in PBS or TBS to yield a 0.1 mg/ml solution. Method 2 for immunoprecipitation, supershift, immunostaining and immunoblotting (Western blot): Pellet antibodies. Remove supernatant. Dissolve the pellet in 50ul of PBS (or TBS) at final concentration of 1 mg/ml. Do not allow the pellet to dry. Do not vortex. Not tested in other applications. Optimal dilutions/concentrations should be determined by the end user. Immunohistochemistry, Western blot.

Antibody Name: NKX3A antibody

Description: This polyclonal targets NKX3A antibody

Target Organism: rat, mouse, human

Antibody ID: AB_372507

Vendor: GeneTex

Catalog Number: GTX12044

Record Creation Time: 20231110T081217+0000

Ratings and Alerts

No rating or validation information has been found for NKX3A antibody.

Warning: Discontinued at GeneTex

Discontinued; manufacturer recommendations: IgG; IgG Immunohistochemistry; Other; Western Blot; IHC: Use at a dilution of 1/200 - 1/1000. WB: Use at a dilution of 1/200 - 1/1000. Detects a band of approximately 21.6 - 26.3 kDa. Preparation InstructionsMethod 1 for immunostaining and immunoblotting (Western blot): Remove a fixed amount of suspension and dissolve 1/5 in PBS or TBS to yield a 0.1 mg/ml solution. Method 2 for immunoprecipitation, supershift, immunostaining and immunoblotting (Western blot): Pellet antibodies. Remove supernatant. Dissolve the pellet in 50ul of PBS (or TBS) at final concentration of 1 mg/ml. Do not allow the pellet to dry. Do not vortex. Not tested in other applications. Optimal dilutions/concentrations should be determined by the end user. Immunohistochemistry, Western blot.

Data and Source Information

Source: [Antibody Registry](#)

Usage and Citation Metrics

We have not found any literature mentions for this resource.