

Resource Summary Report

Generated by [RRID](#) on Jul 30, 2026

PTH antibody

RRID:AB_372670

Type: Antibody

Proper Citation

GeneTex Cat# GTX27843, RRID:AB_372670

Antibody Information

URL: http://antibodyregistry.org/AB_372670

Proper Citation: GeneTex Cat# GTX27843, RRID:AB_372670

Target Antigen: PTH antibody

Host Organism: rabbit

Clonality: polyclonal

Comments: Discontinued; manufacturer recommendations: IgG; IgG Immunohistochemistry; Immunohistochemistry - fixed; Immunofluorescence; Immunohistochemistry - frozen; Immunoprecipitation; Western Blot; For WB: Use at a dilution of 1:100-1:200. For IP: Use at an assay dependent dilution. For IHC: Use at a dilution of 1:40-1:80. This antibody can be used on frozen tissue sections and on formalin-fixed, paraffin-embedded tissue sections. Prolonged fixation in buffered formalin can destroy the epitope. The antibody may be used at a dilution of 1:40-1:80 with Auto/Probe III. Auto/Zyme (Cat GTX73183) predigestion or high temperature antigenic unmasking (see instructions below) is required with this antibody prior to application of this antibody with formalin-fixed, paraffin-embedded tissue. The optimum dilution factor should be determined by individual lab

Antibody Name: PTH antibody

Description: This polyclonal targets PTH antibody

Target Organism: human

Antibody ID: AB_372670

Vendor: GeneTex

Catalog Number: GTX27843

Record Creation Time: 20250820T033705+0000

Record Last Update: 20250820T164337+0000

Ratings and Alerts

No rating or validation information has been found for PTH antibody.

Warning: Discontinued at GeneTex

Discontinued; manufacturer recommendations: IgG; IgG Immunohistochemistry; Immunohistochemistry - fixed; Immunofluorescence; Immunohistochemistry - frozen; Immunoprecipitation; Western Blot; For WB: Use at a dilution of 1:100-1:200. For IP: Use at an assay dependent dilution. For IHC: Use at a dilution of 1:40-1:80. This antibody can be used on frozen tissue sections and on formalin-fixed, paraffin-embedded tissue sections. Prolonged fixation in buffered formalin can destroy the epitope. The antibody may be used at a dilution of 1:40-1:80 with Auto/Probe III. Auto/Zyme (Cat GTX73183) predigestion or high temperature antigenic unmasking (see instructions below) is required with this antibody prior to application of this antibody with formalin-fixed, paraffin-embedded tissue. The optimum dilution factor should be determined by individual lab

Data and Source Information

Source: [Antibody Registry](#)

Usage and Citation Metrics

We have not found any literature mentions for this resource.