

# Resource Summary Report

Generated by [RRID](#) on Sep 9, 2026

## MCP2 antibody

RRID:AB\_373988

Type: Antibody

### Proper Citation

GeneTex Cat# GTX10392, RRID:AB\_373988

### Antibody Information

**URL:** [http://antibodyregistry.org/AB\\_373988](http://antibodyregistry.org/AB_373988)

**Proper Citation:** GeneTex Cat# GTX10392, RRID:AB\_373988

**Target Antigen:** MCP2 antibody

**Host Organism:** goat

**Clonality:** polyclonal

**Comments:** Discontinued; manufacturer recommendations: IgG; IgG ELISA: Use at a concentration of 0.5 - 1 µg/ml to detect approximately 0.3 ng/well of recombinant human MCP2. Neut: Use at a concentration of 2 - 5 µg/ml. WB: Use at a concentration of 0.1 - 0.2 µg/ml. The detection limit is approximately 20ng/lane under non-reducing and reducing conditions. Predicted molecular weight: 13kDa. Neutralization Dose 50 (ND50) is defined as the concentration of antibody required to yield one-half maximal inhibition of the cytokine activity on a responsive cell line, when that cytokine is present at a concentration just high enough to elicit a maximum response. The ND50 is approximately 2 - 5 µg/ml in the presence of 0.4 µg/ml of human MCP2, using the human monocyte chemotaxis assay. Not tested in other applications. Optimal dilutio

**Antibody Name:** MCP2 antibody

**Description:** This polyclonal targets MCP2 antibody

**Target Organism:** human

**Antibody ID:** AB\_373988

**Vendor:** GeneTex

**Catalog Number:** GTX10392

**Record Creation Time:** 20231110T081216+0000

## Ratings and Alerts

No rating or validation information has been found for MCP2 antibody.

**Warning:** Discontinued at GeneTex

Discontinued; manufacturer recommendations: IgG; IgG ELISA: Use at a concentration of 0.5 - 1 µg/ml to detect approximately 0.3 ng/well of recombinant human MCP2. Neut: Use at a concentration of 2 - 5 µg/ml. WB: Use at a concentration of 0.1 - 0.2 µg/ml. The detection limit is approximately 20ng/lane under non-reducing and reducing conditions. Predicted molecular weight: 13kDa. Neutralization Dose 50 (ND50) is defined as the concentration of antibody required to yield one-half maximal inhibition of the cytokine activity on a responsive cell line, when that cytokine is present at a concentration just high enough to elicit a maximum response. The ND50 is approximately 2 - 5 µg/ml in the presence of 0.4 µg/ml of human MCP2, using the human monocyte chemotaxis assay. Not tested in other applications. Optimal dilutio

---

## Data and Source Information

**Source:** [Antibody Registry](#)

---

## Usage and Citation Metrics

We have not found any literature mentions for this resource.