

Resource Summary Report

Generated by [RRID](#) on Aug 5, 2026

lambda Light chain [HP6054] antibody

RRID:AB_374989

Type: Antibody

Proper Citation

GeneTex Cat# GTX73553, RRID:AB_374989

Antibody Information

URL: http://antibodyregistry.org/AB_374989

Proper Citation: GeneTex Cat# GTX73553, RRID:AB_374989

Target Antigen: lambda Light chain [HP6054] antibody

Host Organism: mouse

Clonality: monoclonal

Comments: Discontinued; manufacturer recommendations: Immunohistochemistry; Immunohistochemistry - fixed; IHC, IHC-P, Immunohistochemistry, IHC (Formalin-fixed paraffin-embedded sections). The usefulness of this product in other applications has not been determined., This antibody may be diluted to a titer of 1:50-1:100 in an ABC method. However, depending on the staining conditions employed, we suggest that final dilution should be determined by the user. We suggest an incubation period of 30 minutes at room temperature. However, depending upon the fixation conditions and the staining system employed, optimal incubation should be determined by the user. Formalin fixed paraffin embedded tissue sections require high temperature antigen unmasking with 10 mM citrate buffer, pH 6.0 prior to immunostaining.

Antibody Name: lambda Light chain [HP6054] antibody

Description: This monoclonal targets lambda Light chain [HP6054] antibody

Antibody ID: AB_374989

Vendor: GeneTex

Catalog Number: GTX73553

Record Creation Time: 20250820T013409+0000

Record Last Update: 20250820T111634+0000

Ratings and Alerts

No rating or validation information has been found for lambda Light chain [HP6054] antibody.

Warning: Discontinued at GeneTex

Discontinued; manufacturer recommendations: Immunohistochemistry; Immunohistochemistry - fixed; IHC, IHC-P, Immunohistochemistry, IHC (Formalin-fixed paraffin-embedded sections). The usefulness of this product in other applications has not been determined., This antibody may be diluted to a titer of 1:50-1:100 in an ABC method. However, depending on the staining conditions employed, we suggest that final dilution should be determined by the user. We suggest an incubation period of 30 minutes at room temperature. However, depending upon the fixation conditions and the staining system employed, optimal incubation should be determined by the user. Formalin fixed paraffin embedded tissue sections require high temperature antigen unmasking with 10 mM citrate buffer, pH 6.0 prior to immunostaining.

Data and Source Information

Source: [Antibody Registry](#)

Usage and Citation Metrics

We have not found any literature mentions for this resource.