

Resource Summary Report

Generated by [RRID](#) on Aug 11, 2026

CD4 [1F6] antibody (ready-to-use)

RRID:AB_375098

Type: Antibody

Proper Citation

GeneTex Cat# GTX73720, RRID:AB_375098

Antibody Information

URL: http://antibodyregistry.org/AB_375098

Proper Citation: GeneTex Cat# GTX73720, RRID:AB_375098

Target Antigen: CD4 [1F6] antibody (ready-to-use)

Host Organism: mouse

Clonality: monoclonal

Comments: Discontinued; manufacturer recommendations: Immunohistochemistry; Immunohistochemistry - fixed; Immunohistochemistry - frozen; IHC, IHC-P, Immunohistochemistry, IHC (Formalin-fixed paraffin-embedded sections). The usefulness of this product in other applications has not been determined., This antibody has been pre-titrated and quality controlled to work on formalin-fixed paraffin-embedded as well as acetone fixed cryostat tissue sections. No further titration is required. We suggest an incubation period of 30 to 60 minutes at room temperature or overnight at 40 C. However, depending upon the fixation conditions and the staining system employed, optimal incubation should be determined by the user. High temperature antigen unmasking technique is required for paraffin embedded tissue sections. We recommend using 1mM EDTA pH 8.0 buffer in

Antibody Name: CD4 [1F6] antibody (ready-to-use)

Description: This monoclonal targets CD4 [1F6] antibody (ready-to-use)

Antibody ID: AB_375098

Vendor: GeneTex

Catalog Number: GTX73720

Record Creation Time: 20250819T215227+0000

Record Last Update: 20250820T002218+0000

Ratings and Alerts

No rating or validation information has been found for CD4 [1F6] antibody (ready-to-use).

Warning: Discontinued at GeneTex

Discontinued; manufacturer recommendations: Immunohistochemistry; Immunohistochemistry - fixed; Immunohistochemistry - frozen; IHC, IHC-P, Immunohistochemistry, IHC (Formalin-fixed paraffin-embedded sections). The usefulness of this product in other applications has not been determined., This antibody has been pretitered and quality controlled to work on formalin-fixed paraffin-embedded as well as acetone fixed cryostat tissue sections. No further titration is required. We suggest an incubation period of 30 to 60 minutes at room temperature or overnight at 40 C. However, depending upon the fixation conditions and the staining system employed, optimal incubation should be determined by the user. High temperature antigen unmasking technique is required for paraffin embedded tissue sections. We recommend using 1mM EDTA pH 8.0 buffer in

Data and Source Information

Source: [Antibody Registry](#)

Usage and Citation Metrics

We have not found any literature mentions for this resource.