

Resource Summary Report

Generated by [RRID](#) on Aug 8, 2026

CD35 [E11] antibody

RRID:AB_376064

Type: Antibody

Proper Citation

GeneTex Cat# GTX73816, RRID:AB_376064

Antibody Information

URL: http://antibodyregistry.org/AB_376064

Proper Citation: GeneTex Cat# GTX73816, RRID:AB_376064

Target Antigen: CD35 [E11] antibody

Host Organism: mouse

Clonality: monoclonal

Comments: Discontinued; manufacturer recommendations: Immunohistochemistry; Immunohistochemistry - fixed; This antibody may be diluted to a titer of 1:25-1:50 in an ABC method. However, depending on the staining conditions employed, we suggest that final dilution should be determined by the user. We suggest an incubation period of 30 minutes at room temperature. However, depending upon the fixation conditions and the staining system employed, optimal incubation should be determined by the user. Formalin fixed paraffin embedded tissue sections require high temperature antigen unmasking with 10 mM citrate buffer, pH 6.0 prior to immunostaining, IHC, IHC-P, Immunohistochemistry, IHC (Formalin-fixed paraffin-embedded sections). The usefulness of this product in other applications has not been determined.

Antibody Name: CD35 [E11] antibody

Description: This monoclonal targets CD35 [E11] antibody

Antibody ID: AB_376064

Vendor: GeneTex

Catalog Number: GTX73816

Record Creation Time: 20250820T055420+0000

Record Last Update: 20250820T224646+0000

Ratings and Alerts

No rating or validation information has been found for CD35 [E11] antibody.

Warning: Discontinued at GeneTex

Discontinued; manufacturer recommendations: Immunohistochemistry; Immunohistochemistry - fixed; This antibody may be diluted to a titer of 1:25-1:50 in an ABC method. However, depending on the staining conditions employed, we suggest that final dilution should be determined by the user. We suggest an incubation period of 30 minutes at room temperature. However, depending upon the fixation conditions and the staining system employed, optimal incubation should be determined by the user. Formalin fixed paraffin embedded tissue sections require high temperature antigen unmasking with 10 mM citrate buffer, pH 6.0 prior to immunostaining, IHC, IHC-P, Immunohistochemistry, IHC (Formalin-fixed paraffin-embedded sections). The usefulness of this product in other applications has not been determined.

Data and Source Information

Source: [Antibody Registry](#)

Usage and Citation Metrics

We have not found any literature mentions for this resource.