

Resource Summary Report

Generated by [RRID](#) on Sep 10, 2026

HSV1 [20.7.1] antibody (ready-to-use)

RRID:AB_379512

Type: Antibody

Proper Citation

GeneTex Cat# GTX73490, RRID:AB_379512

Antibody Information

URL: http://antibodyregistry.org/AB_379512

Proper Citation: GeneTex Cat# GTX73490, RRID:AB_379512

Target Antigen: HSV1 [20.7.1] antibody (ready-to-use)

Host Organism: mouse

Clonality: monoclonal

Comments: Discontinued; manufacturer recommendations: This antibody has been pretitered and quality controlled to work on formalin-fixed paraffin-embedded as well as acetone fixed cryostat tissue sections. No further titration is required. We suggest an incubation period of 30 minutes at room temperature. However, depending upon the fixation conditions and the staining system employed, optimal incubation should be determined by the user. Proteolytic treatment is required prior to immunostaining of formalin fixed paraffin embedded tissue sections., IHC, IHC-P, Immunohistochemistry, IHC (Formalin-fixed paraffin-embedded sections). The usefulness of this product in other applications has not been determined.; Immunohistochemistry - frozen; Immunohistochemistry; Immunohistochemistry - fixed

Antibody Name: HSV1 [20.7.1] antibody (ready-to-use)

Description: This monoclonal targets HSV1 [20.7.1] antibody (ready-to-use)

Antibody ID: AB_379512

Vendor: GeneTex

Catalog Number: GTX73490

Record Creation Time: 20231110T081226+0000

Record Last Update: 20260829T063911+0000

Ratings and Alerts

No rating or validation information has been found for HSV1 [20.7.1] antibody (ready-to-use).

Warning: Discontinued at GeneTex

Discontinued; manufacturer recommendations: This antibody has been pretitered and quality controlled to work on formalin-fixed paraffin-embedded as well as acetone fixed cryostat tissue sections. No further titration is required. We suggest an incubation period of 30 minutes at room temperature. However, depending upon the fixation conditions and the staining system employed, optimal incubation should be determined by the user. Proteolytic treatment is required prior to immunostaining of formalin fixed paraffin embedded tissue sections., IHC, IHC-P, Immunohistochemistry, IHC (Formalin-fixed paraffin-embedded sections). The usefulness of this product in other applications has not been determined.; Immunohistochemistry - frozen; Immunohistochemistry; Immunohistochemistry - fixed

Data and Source Information

Source: [Antibody Registry](#)

Usage and Citation Metrics

We have not found any literature mentions for this resource.