

Resource Summary Report

Generated by [RRID](#) on Aug 10, 2026

Apo E [E6D7] antibody

RRID:AB_380317

Type: Antibody

Proper Citation

GeneTex Cat# GTX21907, RRID:AB_380317

Antibody Information

URL: http://antibodyregistry.org/AB_380317

Proper Citation: GeneTex Cat# GTX21907, RRID:AB_380317

Target Antigen: Apo E [E6D7] antibody

Host Organism: mouse

Clonality: monoclonal

Comments: Discontinued; manufacturer recommendations: Immunohistochemistry - frozen; Western Blot; Immunohistochemistry - fixed; ELISA; Immunohistochemistry; Immunoprecipitation; ELISA, IHC (Frozen sections), IHC (Formalin-fixed paraffin-embedded sections), Immunoprecipitation, Western blot. The usefulness of this product in other applications has not been determined., ELISA, IHC-Fr, IHC-P, IP, WB, ELISA: Use at an assay dependent dilution. IHC-P: Use at an assay dependent dilution. Staining may be significantly improved by either1. Formic acid (70 %) for 10-30 minutes at room temperature2. HIER (heat Induced Epitope Retrieval) with buffers such as Citrate IHC-Fr: Use at an assay dependent dilution. IP: Use at an assay dependent dilution. WB: Use at a dilution of 1/100 - 1/1000. Predicted molecular weight: 38 kDa. Not tested in other applicati

Antibody Name: Apo E [E6D7] antibody

Description: This monoclonal targets Apo E [E6D7] antibody

Antibody ID: AB_380317

Vendor: GeneTex

Catalog Number: GTX21907

Record Creation Time: 20250820T000123+0000

Record Last Update: 20250820T070841+0000

Ratings and Alerts

No rating or validation information has been found for Apo E [E6D7] antibody.

Warning: Discontinued at GeneTex

Discontinued; manufacturer recommendations: Immunohistochemistry - frozen; Western Blot; Immunohistochemistry - fixed; ELISA; Immunohistochemistry; Immunoprecipitation; ELISA, IHC (Frozen sections), IHC (Formalin-fixed paraffin-embedded sections), Immunoprecipitation, Western blot. The usefulness of this product in other applications has not been determined., ELISA, IHC-Fr, IHC-P, IP, WB, ELISA: Use at an assay dependent dilution. IHC-P: Use at an assay dependent dilution. Staining may be significantly improved by either1. Formic acid (70) for 10-30 minutes at room temperature2. HIER (heat Induced Epitope Retrieval) with buffers such as Citrate IHC-Fr: Use at an assay dependent dilution. IP: Use at an assay dependent dilution. WB: Use at a dilution of 1/100 - 1/1000. Predicted molecular weight: 38 kDa. Not tested in other applicati

Data and Source Information

Source: [Antibody Registry](#)

Usage and Citation Metrics

We have not found any literature mentions for this resource.