

Resource Summary Report

Generated by [RRID](#) on Jul 28, 2026

Interferon gamma antibody

RRID:AB_381144

Type: Antibody

Proper Citation

GeneTex Cat# GTX10744, RRID:AB_381144

Antibody Information

URL: http://antibodyregistry.org/AB_381144

Proper Citation: GeneTex Cat# GTX10744, RRID:AB_381144

Target Antigen: Interferon gamma antibody

Host Organism: goat

Clonality: polyclonal

Comments: Discontinued; manufacturer recommendations: IgG; IgG Block/Neutralize/Inhibit; Immunohistochemistry; ELISA; Western Blot; Other; ELISA, Immunohistochemistry, Neutralizing, Western blot. ELISA: Use at a concentration of 0.5 - 1 µg/ml. The detection limit for recombinant rat IFNG and recombinant mouse IFNG is approximately 0.2 ng/well. IHC: Use at a concentration of 0.5 - 5 µg/ml this concentration is recommended to detect rat IFNG in cultured cells or tissue sections. Neut: Use at a concentration of 0.08 - 0.2 µg/ml. WB: Use at a concentration of 0.1 - 0.2 µg/ml. The detection limit for recombinant rat IFNG and recombinant mouse IFNG is approximately 5 ng/lane under non-reducing and reducing conditions. Predicted molecular weight: 15.5 kDa. To measure the ability of this antibody to neutralize the bioactivity of rat I

Antibody Name: Interferon gamma antibody

Description: This polyclonal targets Interferon gamma antibody

Target Organism: rat, mouse

Antibody ID: AB_381144

Vendor: GeneTex

Catalog Number: GTX10744

Record Creation Time: 20250820T040159+0000

Ratings and Alerts

No rating or validation information has been found for Interferon gamma antibody.

Warning: Discontinued at GeneTex

Discontinued; manufacturer recommendations: IgG; IgG Block/Neutralize/Inhibit; Immunohistochemistry; ELISA; Western Blot; Other; ELISA, Immunohistochemistry, Neutralizing, Western blot. ELISA: Use at a concentration of 0.5 - 1 µg/ml. The detection limit for recombinant rat IFNG and recombinant mouse IFNG is approximately 0.2 ng/wellIHC: Use at a concentration of 0.5 - 5 µg/ml this concentration is recommended to detect rat IFNG in cultured cells or tissue sections. Neut: Use at a concentration of 0.08 - 0.2 µg/ml. WB: Use at a concentration of 0.1 - 0.2 µg/ml. The detection limit for recombinant rat IFNG and recombinant mouse IFNG is approximately 5 ng/lane under non-reducing and reducing conditions. Predicted molecular weight: 15.5 kDa. To measure the ability of this antibody to neutralize the bioactivity of rat I

Data and Source Information

Source: [Antibody Registry](#)

Usage and Citation Metrics

We have not found any literature mentions for this resource.