

Resource Summary Report

Generated by [RRID](#) on Sep 10, 2026

VEGF receptor 2 antibody [KDR-2 or 260.4] (Biotin)

RRID:AB_381210

Type: Antibody

Proper Citation

GeneTex Cat# GTX10975, RRID:AB_381210

Antibody Information

URL: http://antibodyregistry.org/AB_381210

Proper Citation: GeneTex Cat# GTX10975, RRID:AB_381210

Target Antigen: VEGF receptor 2 antibody [KDR-2 or 260.4] ()

Host Organism: mouse

Clonality: monoclonal

Comments: Discontinued; manufacturer recommendations: IgG1; IgG1 Flow Cytometry; Western Blot; Immunofluorescence; Immunohistochemistry; ELISA; ELISA, Flow cytometry/FACS, Immunohistochemistry, Western blot. The usefulness of this product in other applications has not been determined., ELISA, FACS, IHC, WB, ELISA: Use at an assay dependent dilution. FACS: Use at an assay dependent dilution. IHC: Use at an assay dependent dilution. WB: Use at a dilution of 1/1000. This concentration is determined using a culture supernatant of insect cells infected with a baculovirus expressing recombinant extracellular human KDR receptor. Predicted molecular weight: 152 kDa. Not tested in other applications. Optimal dilutions/concentrations should be determined by the end user.

Antibody Name: VEGF receptor 2 antibody [KDR-2 or 260.4] (Biotin)

Description: This monoclonal targets VEGF receptor 2 antibody [KDR-2 or 260.4] ()

Antibody ID: AB_381210

Vendor: GeneTex

Catalog Number: GTX10975

Record Creation Time: 20231110T081054+0000

Record Last Update: 20260829T114957+0000

Ratings and Alerts

No rating or validation information has been found for VEGF receptor 2 antibody [KDR-2 or 260.4] (Biotin).

Warning: Discontinued at GeneTex

Discontinued; manufacturer recommendations: IgG1; IgG1 Flow Cytometry; Western Blot; Immunofluorescence; Immunohistochemistry; ELISA; ELISA, Flow cytometry/FACS, Immunohistochemistry, Western blot. The usefulness of this product in other applications has not been determined., ELISA, FACS, IHC, WB, ELISA: Use at an assay dependent dilution. FACS: Use at an assay dependent dilution. IHC: Use at an assay dependent dilution. WB: Use at a dilution of 1/1000. This concentration is determined using a culture supernatant of insect cells infected with a baculovirus expressing recombinant extracellular human KDR receptor. Predicted molecular weight: 152 kDa. Not tested in other applications. Optimal dilutions/concentrations should be determined by the end user.

Data and Source Information

Source: [Antibody Registry](#)

Usage and Citation Metrics

We have not found any literature mentions for this resource.