

Resource Summary Report

Generated by [RRID](#) on Aug 9, 2026

TIMP4 antibody

RRID:AB_383782

Type: Antibody

Proper Citation

GeneTex Cat# GTX73964, RRID:AB_383782

Antibody Information

URL: http://antibodyregistry.org/AB_383782

Proper Citation: GeneTex Cat# GTX73964, RRID:AB_383782

Target Antigen: TIMP4 antibody

Host Organism: rabbit

Clonality: polyclonal

Comments: Discontinued; manufacturer recommendations: IgG; IgG Immunohistochemistry; Immunohistochemistry - fixed; Immunohistochemistry, IHC (Formalin-fixed paraffin-embedded sections). This antibody may be diluted to a titer of 1:25-1:50 in an ABC method. However, depending on the staining conditions employed, we suggest that final dilution should be determined by the user. We suggest an incubation period of 30 minutes at room temperature. However, depending upon the fixation conditions and the staining system employed, optimal incubation conditions should be determined by the user. Formalin fixed paraffin embedded tissue sections require high temperature antigen unmasking with 10 mM citrate buffer, pH 6.0 prior to immunostaining.

Antibody Name: TIMP4 antibody

Description: This polyclonal targets TIMP4 antibody

Target Organism: human

Antibody ID: AB_383782

Vendor: GeneTex

Catalog Number: GTX73964

Record Creation Time: 20250820T002537+0000

Record Last Update: 20250820T081419+0000

Ratings and Alerts

No rating or validation information has been found for TIMP4 antibody.

Warning: Discontinued at GeneTex

Discontinued; manufacturer recommendations: IgG; IgG Immunohistochemistry; Immunohistochemistry - fixed; Immunohistochemistry, IHC (Formalin-fixed paraffin-embedded sections). This antibody may be diluted to a titer of 1:25-1:50 in an ABC method. However, depending on the staining conditions employed, we suggest that final dilution should be determined by the user. We suggest an incubation period of 30 minutes at room temperature. However, depending upon the fixation conditions and the staining system employed, optimal incubation conditions should be determined by the user. Formalin fixed paraffin embedded tissue sections require high temperature antigen unmasking with 10 mM citrate buffer, pH 6.0 prior to immunostaining.

Data and Source Information

Source: [Antibody Registry](#)

Usage and Citation Metrics

We have not found any literature mentions for this resource.