

Resource Summary Report

Generated by [RRID](#) on Sep 11, 2026

cN RAS [F155-227] antibody

RRID:AB_384288

Type: Antibody

Proper Citation

GeneTex Cat# GTX13054, RRID:AB_384288

Antibody Information

URL: http://antibodyregistry.org/AB_384288

Proper Citation: GeneTex Cat# GTX13054, RRID:AB_384288

Target Antigen: cN RAS [F155-227] antibody

Host Organism: mouse

Clonality: monoclonal

Comments: Discontinued; manufacturer recommendations: Immunoprecipitation; Immunohistochemistry; Immunohistochemistry - frozen; Immunohistochemistry - fixed; Western Blot; For IHC-P: Use at a concentration of 8 - 12 µg/ml. Anti-trp E may be used as a negative control. Paraffin sections will require treatment with saponin (0.05 in water, 30 min.) or pepsin (0.1 in 0.1N HCl, 10-20 min.) at room temperature. For IHC-Fr: Use at a concentration of 8 - 12 µg/ml. For IP: Use 5 µg per sample with 45 µl of Protein G-agarose. For WB: Use at a concentration of 1-10µg/ml. This concentration is recommended using a neuronal cell extract. Predicted molecular weight: 21 kDa. Not tested in other applications. Optimal dilutions/concentrations should be determined by the researcher., IHC-Fr, IHC-P, IP, WB, IHC (Frozen sections),

Antibody Name: cN RAS [F155-227] antibody

Description: This monoclonal targets cN RAS [F155-227] antibody

Target Organism: rat, mouse, human

Antibody ID: AB_384288

Vendor: GeneTex

Catalog Number: GTX13054

Record Creation Time: 20231110T081056+0000

Ratings and Alerts

No rating or validation information has been found for cN RAS [F155-227] antibody.

Warning: Discontinued at GeneTex

Discontinued; manufacturer recommendations: Immunoprecipitation; Immunohistochemistry; Immunohistochemistry - frozen; Immunohistochemistry - fixed; Western Blot; For IHC-P: Use at a concentration of 8 - 12 µg/ml. Anti-trp E may be used as a negative control. Paraffin sections will require treatment with saponin (0.05 in water, 30 min.) or pepsin (0.1 in 0.1N HCl, 10-20 min.) at room temperature. For IHC-Fr: Use at a concentration of 8 - 12 µg/ml. For IP: Use 5 µg per sample with 45 µl of Protein G-agarose. For WB: Use at a concentration of 1-10µg/ml. This concentration is recommended using a neuronal cell extract. Predicted molecular weight: 21 kDa. Not tested in other applications. Optimal dilutions/concentrations should be determined by the researcher., IHC-Fr, IHC-P, IP, WB, IHC (Frozen sections),

Data and Source Information

Source: [Antibody Registry](#)

Usage and Citation Metrics

We have not found any literature mentions for this resource.