

Resource Summary Report

Generated by [RRID](#) on Aug 5, 2026

Hairy Cell Leukemia [DBA.44] antibody

RRID:AB_384480

Type: Antibody

Proper Citation

GeneTex Cat# GTX20779, RRID:AB_384480

Antibody Information

URL: http://antibodyregistry.org/AB_384480

Proper Citation: GeneTex Cat# GTX20779, RRID:AB_384480

Target Antigen: Hairy Cell Leukemia [DBA.44] antibody

Host Organism: mouse

Clonality: monoclonal

Comments: Discontinued; manufacturer recommendations: For IHC: This antibody may be diluted to a titer of 1:15-1:30 in an ABC method. Staining Protocol: We suggest an incubation period of 60 minutes at room temperature. However, depending upon the fixation conditions and the staining system employed, optimal incubation conditions and antibody dilutions should be determined by the user. Enzymatic antigen retrieval of formalin fixed paraffin allembded tissue sections is required prior to immunostaining. Not tested in other applications. Optimal dilutions/concentrations should be determined by the researcher., IHC (Frozen sections), IHC (Formalin-fixed paraffin-embedded sections). The usefulness of this product in other applications has not been determined., IHC-Fr, IHC-P; Immunohistochemistry - frozen; Immunohistochemistry; Immunohistochemi

Antibody Name: Hairy Cell Leukemia [DBA.44] antibody

Description: This monoclonal targets Hairy Cell Leukemia [DBA.44] antibody

Antibody ID: AB_384480

Vendor: GeneTex

Catalog Number: GTX20779

Record Creation Time: 20250820T055721+0000

Record Last Update: 20250820T225445+0000

Ratings and Alerts

No rating or validation information has been found for Hairy Cell Leukemia [DBA.44] antibody.

Warning: Discontinued at GeneTex

Discontinued; manufacturer recommendations: For IHC: This antibody may be diluted to a titer of 1:15-1:30 in an ABC method. Staining Protocol: We suggest an incubation period of 60 minutes at room temperature. However, depending upon the fixation conditions and the staining system employed, optimal incubation conditions and antibody dilutions should be determined by the user. Enzymatic antigen retrieval of formalin fixed paraffin allembded tissue sections is required prior to immunostaining. Not tested in other applications. Optimal dilutions/concentrations should be determined by the researcher., IHC (Frozen sections), IHC (Formalin-fixed paraffin-embedded sections). The usefulness of this product in other applications has not been determined., IHC-Fr, IHC-P; Immunohistochemistry - frozen; Immunohistochemistry; Immunohistochemi

Data and Source Information

Source: [Antibody Registry](#)

Usage and Citation Metrics

We have not found any literature mentions for this resource.