

Resource Summary Report

Generated by [RRID](#) on Aug 4, 2026

Hydrogen/Potassium ATPase beta antibody [2G11]

RRID:AB_384885

Type: Antibody

Proper Citation

GeneTex Cat# GTX22866, RRID:AB_384885

Antibody Information

URL: http://antibodyregistry.org/AB_384885

Proper Citation: GeneTex Cat# GTX22866, RRID:AB_384885

Target Antigen: Hydrogen/Potassium ATPase beta antibody [2G11]

Host Organism: mouse

Clonality: monoclonal

Comments: Discontinued; manufacturer recommendations: IgG1; IgG1 ICC, IF, IHC, IHC-P, IP, WB, Immunocytochemistry, Immunofluorescence, Immunohistochemistry, IHC (Formalin-fixed paraffin-embedded sections), Immunoprecipitation, Western Blot. The usefulness of this product in other applications has not been determined., For IHC: use at a dilution of 1/2000 For WB: use at a dilution of 1/4000Not tested in other applications. Optimal dilutions / concentrations should be determined by the end user. By Western blot, this antibody detects various forms of the beta-subunit, including a 60 to 80 kDa glycosylated protein, a 52 kDa beta-subunit precursor, and the 34 kDa core peptide from rabbit stomach extracts. Immunohistochemical staining of hydrogen/potassium ATPase in rabbit gastric glands with this antibody results in specific staining of parietal cells. This

Antibody Name: Hydrogen/Potassium ATPase beta antibody [2G11]

Description: This monoclonal targets Hydrogen/Potassium ATPase beta antibody [2G11]

Target Organism: rat, porcine, canine, pig, mouse, rabbit, bovine, dog, human

Antibody ID: AB_384885

Vendor: GeneTex

Catalog Number: GTX22866

Record Creation Time: 20250820T012014+0000

Ratings and Alerts

No rating or validation information has been found for Hydrogen/Potassium ATPase beta antibody [2G11].

Warning: Discontinued at GeneTex

Discontinued; manufacturer recommendations: IgG1; IgG1 ICC, IF, IHC, IHC-P, IP, WB, Immunocytochemistry, Immunofluorescence, Immunohistochemistry, IHC (Formalin-fixed paraffin-embedded sections), Immunoprecipitation, Western Blot. The usefulness of this product in other applications has not been determined., For IHC: use at a dilution of 1/2000 For WB: use at a dilution of 1/4000Not tested in other applications. Optimal dilutions / concentrations should be determined by the end user. By Western blot, this antibody detects various forms of the beta-subunit, including a 60 to 80 kDa glycosylated protein, a 52 kDa beta-subunit precursor, and the 34 kDa core peptide from rabbit stomach extracts. Immunohistochemical staining of hydrogen/potassium ATPase in rabbit gastric glands with this antibody results in specific staining of parietal cells. This

Data and Source Information

Source: [Antibody Registry](#)

Usage and Citation Metrics

We have not found any literature mentions for this resource.