

Resource Summary Report

Generated by [RRID](#) on Sep 9, 2026

P Glycoprotein [265/F4] antibody

RRID:AB_384932

Type: Antibody

Proper Citation

GeneTex Cat# GTX23083, RRID:AB_384932

Antibody Information

URL: http://antibodyregistry.org/AB_384932

Proper Citation: GeneTex Cat# GTX23083, RRID:AB_384932

Target Antigen: P Glycoprotein [265/F4] antibody

Host Organism: mouse

Clonality: monoclonal

Comments: Discontinued; manufacturer recommendations: Immunohistochemistry; Western Blot; Immunofluorescence; Flow Cytometry; Immunohistochemistry - frozen; FACS, IF, IHC-Fr, WB, Flow cytometry/FACS, Immunofluorescence, IHC (Frozen sections), Western blot. The usefulness of this product in other applications has not been determined., Flow cytometry / FACS: assay dependent. Immunofluorescence: assay dependent. Western blot: use at a concentration of 1 - 2µg/ml for 2 hours at RT. Immunohistochemistry (frozen sections): assay dependent. Not tested in other applications. Optimal dilutions / concentrations should be determined by the end user. By Western blot, this antibody detects a band of ~ 170 kDa, which corresponds to the predicted molecular weight of P glycoprotein.

Antibody Name: P Glycoprotein [265/F4] antibody

Description: This monoclonal targets P Glycoprotein [265/F4] antibody

Target Organism: hamster, human

Antibody ID: AB_384932

Vendor: GeneTex

Catalog Number: GTX23083

Record Creation Time: 20231110T081208+0000

Ratings and Alerts

No rating or validation information has been found for P Glycoprotein [265/F4] antibody.

Warning: Discontinued at GeneTex

Discontinued; manufacturer recommendations: Immunohistochemistry; Western Blot; Immunofluorescence; Flow Cytometry; Immunohistochemistry - frozen; FACS, IF, IHC-Fr, WB, Flow cytometry/FACS, Immunofluorescence, IHC (Frozen sections), Western blot. The usefulness of this product in other applications has not been determined., Flow cytometry / FACS: assay dependent. Immunofluorescence: assay dependent. Western blot: use at a concentration of 1 - 2µg/ml for 2 hours at RT. Immunohistochemistry (frozen sections):assay dependent. Not tested in other applications. Optimal dilutions / concentrations should be determined by the end user. By Western blot, this antibody detects a band of ~ 170 kDa, which corresponds to the predicted molecular weight of P glycoprotein.

Data and Source Information

Source: [Antibody Registry](#)

Usage and Citation Metrics

We have not found any literature mentions for this resource.