

Resource Summary Report

Generated by [RRID](#) on Sep 7, 2026

Cytokeratin 5 + 6 [D5/16B4] antibody (ready-to-use)

RRID:AB_422549

Type: Antibody

Proper Citation

GeneTex Cat# GTX17068, RRID:AB_422549

Antibody Information

URL: http://antibodyregistry.org/AB_422549

Proper Citation: GeneTex Cat# GTX17068, RRID:AB_422549

Target Antigen: Cytokeratin 5 + 6 [D5/16B4] antibody (ready-to-use)

Host Organism: mouse

Clonality: monoclonal

Comments: Discontinued; manufacturer recommendations: Immunohistochemistry; Immunohistochemistry - frozen; Immunohistochemistry - fixed; IHC-F, IHC-Fr, IHC-P, IHC (Formalin fixed sections), IHC (Frozen sections), IHC (Formalin-fixed paraffin-embedded sections). The usefulness of this product in other applications has not been determined., IHC-P: Prediluted and ready to use. We suggest an incubation of 30 minutes. Perform heat mediated antigen retrieval before commencing with IHC staining protocol. IHC-Fr: Prediluted and ready to use on acetone fixed cryostat sections. We suggest an incubation of 30 minutes. Not tested in other applications. Optimal dilutions/concentrations should be determined by the end user.

Antibody Name: Cytokeratin 5 + 6 [D5/16B4] antibody (ready-to-use)

Description: This monoclonal targets Cytokeratin 5 + 6 [D5/16B4] antibody (ready-to-use)

Antibody ID: AB_422549

Vendor: GeneTex

Catalog Number: GTX17068

Record Creation Time: 20231110T081055+0000

Record Last Update: 20260901T035523+0000

Ratings and Alerts

No rating or validation information has been found for Cytokeratin 5 + 6 [D5/16B4] antibody (ready-to-use).

Warning: Discontinued at GeneTex

Discontinued; manufacturer recommendations: Immunohistochemistry; Immunohistochemistry - frozen; Immunohistochemistry - fixed; IHC-F, IHC-Fr, IHC-P, IHC (Formalin fixed sections), IHC (Frozen sections), IHC (Formalin-fixed paraffin-embedded sections). The usefulness of this product in other applications has not been determined., IHC-P: Prediluted and ready to use. We suggest an incubation of 30 minutes. Perform heat mediated antigen retrieval before commencing with IHC staining protocol. IHC-Fr: Prediluted and ready to use on acetone fixed cryostat sections. We suggest an incubation of 30 minutes. Not tested in other applications. Optimal dilutions/concentrations should be determined by the end user.

Data and Source Information

Source: [Antibody Registry](#)

Usage and Citation Metrics

We have not found any literature mentions for this resource.