

Resource Summary Report

Generated by [RRID](#) on Sep 8, 2026

p53 [3F301] antibody

RRID:AB_422928

Type: Antibody

Proper Citation

GeneTex Cat# GTX17869, RRID:AB_422928

Antibody Information

URL: http://antibodyregistry.org/AB_422928

Proper Citation: GeneTex Cat# GTX17869, RRID:AB_422928

Target Antigen: p53 [3F301] antibody

Host Organism: mouse

Clonality: monoclonal

Comments: Discontinued; manufacturer recommendations: Immunohistochemistry; Immunohistochemistry - fixed; Immunofluorescence; Flow Cytometry; Western Blot; Immunoprecipitation; FACS: Use at an assay dependent dilution. IHC-P: Use at a concentration of 2 - 4 µg/ml. Staining of formalin/paraffin tissues requires boiling tissue sections in 10mM citrate buffer, pH 6.0, for 10-20 min followed by cooling at RT for 20 min. IP: Use at a concentration of 2 µg/mg lysate (GTX17869 will detect both native and denatured forms of the target, use protein G beads). WB: Use at a concentration of 0.5 - 1 µg/ml. Detects a band of approximately 53 kDa. Not tested in other applications. Optimal dilutions/concentrations should be determined by the end user., FACS, IHC-P, IP, WB, Flow cytometry/FACS, IHC (Formalin-fixed paraffin-embedded sections),

Antibody Name: p53 [3F301] antibody

Description: This monoclonal targets p53 [3F301] antibody

Antibody ID: AB_422928

Vendor: GeneTex

Catalog Number: GTX17869

Record Creation Time: 20231110T081024+0000

Record Last Update: 20260829T034720+0000

Ratings and Alerts

No rating or validation information has been found for p53 [3F301] antibody.

Warning: Discontinued at GeneTex

Discontinued; manufacturer recommendations: Immunohistochemistry; Immunohistochemistry - fixed; Immunofluorescence; Flow Cytometry; Western Blot; Immunoprecipitation; FACS: Use at an assay dependent dilution. IHC-P: Use at a concentration of 2 - 4 µg/ml. Staining of formalin/paraffin tissues requires boiling tissue sections in 10mM citrate buffer, pH 6.0, for 10-20 min followed by cooling at RT for 20 min. IP: Use at a concentration of 2 µg/mg lysate (GTX17869 will detect both native and denatured forms of the target, use protein G beads). WB: Use at a concentration of 0.5 - 1 µg/ml. Detects a band of approximately 53 kDa. Not tested in other applications. Optimal dilutions/concentrations should be determined by the end user., FACS, IHC-P, IP, WB, Flow cytometry/FACS, IHC (Formalin-fixed paraffin-embedded sections),

Data and Source Information

Source: [Antibody Registry](#)

Usage and Citation Metrics

We have not found any literature mentions for this resource.