

Resource Summary Report

Generated by [RRID](#) on Sep 8, 2026

Tropomyosin [SPM224] antibody (ready-to-use)

RRID:AB_425157

Type: Antibody

Proper Citation

GeneTex Cat# GTX17796, RRID:AB_425157

Antibody Information

URL: http://antibodyregistry.org/AB_425157

Proper Citation: GeneTex Cat# GTX17796, RRID:AB_425157

Target Antigen: Tropomyosin [SPM224] antibody (ready-to-use)

Host Organism: mouse

Clonality: monoclonal

Comments: Discontinued; manufacturer recommendations: Immunohistochemistry; Immunohistochemistry - fixed; IHC, IHC-P, Immunohistochemistry, IHC (Formalin-fixed paraffin-embedded sections). The usefulness of this product in other applications has not been determined., IHC-P: Use at an assay dependent dilution. Perform heat mediated antigen retrieval before commencing with IHC staining protocol. Boil tissue sections in 10 mM citrate buffer, pH 6.0 for 10 minutes followed by cooling at room temperature for 20 minutes. Not tested in other applications. Optimal dilutions/concentrations should be determined by the end user.

Antibody Name: Tropomyosin [SPM224] antibody (ready-to-use)

Description: This monoclonal targets Tropomyosin [SPM224] antibody (ready-to-use)

Target Organism: rat, chickenbird, human

Antibody ID: AB_425157

Vendor: GeneTex

Catalog Number: GTX17796

Record Creation Time: 20231110T081047+0000

Record Last Update: 20260830T021103+0000

Ratings and Alerts

No rating or validation information has been found for Tropomyosin [SPM224] antibody (ready-to-use).

Warning: Discontinued at GeneTex

Discontinued; manufacturer recommendations: Immunohistochemistry; Immunohistochemistry - fixed; IHC, IHC-P, Immunohistochemistry, IHC (Formalin-fixed paraffin-embedded sections). The usefulness of this product in other applications has not been determined., IHC-P: Use at an assay dependent dilution. Perform heat mediated antigen retrieval before commencing with IHC staining protocol. Boil tissue sections in 10 mM citrate buffer, pH 6.0 for 10 minutes followed by cooling at room temperature for 20 minutes. Not tested in other applications. Optimal dilutions/concentrations should be determined by the end user.

Data and Source Information

Source: [Antibody Registry](#)

Usage and Citation Metrics

We have not found any literature mentions for this resource.