

Resource Summary Report

Generated by [RRID](#) on Sep 9, 2026

MCP1 [S14] antibody

RRID:AB_425178

Type: Antibody

Proper Citation

GeneTex Cat# GTX18677, RRID:AB_425178

Antibody Information

URL: http://antibodyregistry.org/AB_425178

Proper Citation: GeneTex Cat# GTX18677, RRID:AB_425178

Target Antigen: MCP1 [S14] antibody

Host Organism: mouse

Clonality: monoclonal

Comments: Discontinued; manufacturer recommendations: For ELISA: Use at an assay dependent dilution. This antibody can be used as a capture antibody in sandwich ELISA applications for human MCP-1 detection in combination with a monoclonal tracer/detection antibody (HRP-conjugated or biotin-conjugated). Suggested Capture Coating Dose: 0.3µg/well; Substrate: TMB. If the above suggested conditions are followed approximately 1.5 pg/mL of MCP-1 in serum/plasma or 4 pg/mL of MCP-1 in medium can be detected with an assay range of 0-1600 pg/mL. For Neut: Use at a concentration of 0.5 µg/ml. Tests on the ability to inhibit monocyte chemotaxis toward 1 nM recombinant human MCP-1 in blindwell chambers showed that the antibodies were particularly effective at blocking MCP-1 activity at a concentration of 0.5µg/mL. It was also f

Antibody Name: MCP1 [S14] antibody

Description: This monoclonal targets MCP1 [S14] antibody

Antibody ID: AB_425178

Vendor: GeneTex

Catalog Number: GTX18677

Record Creation Time: 20231110T081053+0000

Record Last Update: 20260901T020541+0000

Ratings and Alerts

No rating or validation information has been found for MCP1 [S14] antibody.

Warning: Discontinued at GeneTex

Discontinued; manufacturer recommendations: For ELISA: Use at an assay dependent dilution. This antibody can be used as a capture antibody in sandwich ELISA applications for human MCP-1 detection in combination with a monoclonal tracer/detection antibody (HRP-conjugated or biotin-conjugated). Suggested Capture Coating Dose: 0.3µg/well; Substrate: TMB. If the above suggested conditions are followed approximately 1.5 pg/mL of MCP-1 in serum/plasma or 4 pg/mL of MCP-1 in medium can be detected with an assay range of 0-1600 pg/mL. For Neut: Use at a concentration of 0.5 µg/ml. Tests on the ability to inhibit monocyte chemotaxis toward 1 nM recombinant human MCP-1 in blindwell chambers showed that the antibodies were particularly effective at blocking MCP-1 activity at a concentration of 0.5µg/mL. It was also f

Data and Source Information

Source: [Antibody Registry](#)

Usage and Citation Metrics

We have not found any literature mentions for this resource.