

Resource Summary Report

Generated by [RRID](#) on Aug 9, 2026

SFPQ antibody [B92]

RRID:AB_510140

Type: Antibody

Proper Citation

GeneTex Cat# GTX11825, RRID:AB_510140

Antibody Information

URL: http://antibodyregistry.org/AB_510140

Proper Citation: GeneTex Cat# GTX11825, RRID:AB_510140

Target Antigen: SFPQ antibody [B92]

Host Organism: mouse

Clonality: monoclonal

Comments: Discontinued; manufacturer recommendations: IgG1; IgG1 Affinity purification, Immunocytochemistry, IHC (Frozen sections), Immunoprecipitation, Radioimmunoassay, Western blot. The usefulness of this product in other applications has not been determined., AP, ICC, IHC-Fr, IP, RIA, WB, For AP: Use at an assay dependent dilution (see Shav-Tal et al reference). For ICC: Use at an assay dependent dilution (fix cells 2 min in 3 paraformaldehyde with 0.5 Triton X-100, and then 20 min in 3 paraformaldehyde) (see Lee et al reference). For IHC-Fr: Use at an assay dependent dilution (4 paraformaldehyde-0.5 Triton X-100 treated tissue). For IP: Use at an assay dependent dilution (see Lee et al reference). For RIA: Use at an assay dependent dilution (see Lee et al reference). WB: WB: Use at a concentration of 2 - 4 µg/ml. Detects a band of app

Antibody Name: SFPQ antibody [B92]

Description: This monoclonal targets SFPQ antibody [B92]

Target Organism: human and mouse - not yet tested in other species, mouse, human

Antibody ID: AB_510140

Vendor: GeneTex

Catalog Number: GTX11825

Record Creation Time: 20250820T002620+0000

Record Last Update: 20250820T081620+0000

Ratings and Alerts

No rating or validation information has been found for SFPQ antibody [B92].

Warning: Discontinued at GeneTex

Discontinued; manufacturer recommendations: IgG1; IgG1 Affinity purification, Immunocytochemistry, IHC (Frozen sections), Immunoprecipitation, Radioimmunoassay, Western blot. The usefulness of this product in other applications has not been determined., AP, ICC, IHC-Fr, IP, RIA, WB, For AP: Use at an assay dependent dilution (see Shav-Tal et al reference). For ICC: Use at an assay dependent dilution (fix cells 2 min in 3 paraformaldehyde with 0.5 Triton X-100, and then 20 min in 3 paraformaldehyde) (see Lee et al reference). For IHC-Fr: Use at an assay dependent dilution (4 paraformaldehyde-0.5 Triton X-100 treated tissue). For IP: Use at an assay dependent dilution (see Lee et al reference). For RIA: Use at an assay dependent dilution (see Lee et al reference). WB: WB: Use at a concentration of 2 - 4 µg/ml. Detects a band of app

Data and Source Information

Source: [Antibody Registry](#)

Usage and Citation Metrics

We have not found any literature mentions for this resource.