

Resource Summary Report

Generated by [RRID](#) on Sep 7, 2026

[alpha Internexin \[1D2\] antibody](#)

RRID:AB_625366

Type: Antibody

Proper Citation

GeneTex Cat# GTX30664, RRID:AB_625366

Antibody Information

URL: http://antibodyregistry.org/AB_625366

Proper Citation: GeneTex Cat# GTX30664, RRID:AB_625366

Target Antigen: alpha Internexin [1D2] antibody

Clonality: monoclonal

Comments: Discontinued; manufacturer recommendations: Immunofluorescence; Immunohistochemistry; Western Blot; IF, IHC-P, WB, Immunofluorescence, IHC (Formalin-fixed paraffin-embedded sections), Western blot. The usefulness of this product in other applications has not been determined., GTX30664 can be used on formalin-fixed cells in tissue culture, cryostat sections, and Western blotting. A 66 kDa band can be seen in Western blotting. This antibody is unusually resistant to aldehyde fixation, so that this antibody is ideal for studies of formalin-fixed paraffin-embedded histological sections. Suggested working dilutions: * Western Blot - 1:10,000 (ECL) Immunohistochemistry** - 1:5,000 Immunofluorescence - 1:500 *The investigator should determine the optimal dilution for a specific application. **ABC or other enzymatic amplif

Antibody Name: alpha Internexin [1D2] antibody

Description: This monoclonal targets alpha Internexin [1D2] antibody

Target Organism: all

Antibody ID: AB_625366

Vendor: GeneTex

Catalog Number: GTX30664

Record Creation Time: 20231110T080417+0000

Record Last Update: 20260828T233031+0000

Ratings and Alerts

No rating or validation information has been found for alpha Internexin [1D2] antibody.

Warning: Discontinued at GeneTex

Discontinued; manufacturer recommendations: Immunofluorescence; Immunohistochemistry; Western Blot; IF, IHC-P, WB, Immunofluorescence, IHC (Formalin-fixed paraffin-embedded sections), Western blot. The usefulness of this product in other applications has not been determined., GTX30664 can be used on formalin-fixed cells in tissue culture, cryostat sections, and Western blotting. A 66 kDa band can be seen in Western blotting. This antibody is unusually resistant to aldehyde fixation, so that this antibody is ideal for studies of formalin-fixed paraffin-embedded histological sections. Suggested working dilutions: * Western Blot - 1:10,000 (ECL) Immunohistochemistry** - 1:5,000 Immunofluorescence - 1:500 *The investigator should determine the optimal dilution for a specific application. **ABC or other enzymatic amplif

Data and Source Information

Source: [Antibody Registry](#)

Usage and Citation Metrics

We have not found any literature mentions for this resource.