

Resource Summary Report

Generated by [RRID](#) on Sep 7, 2026

IgG Light chain [IVA285-1] antibody

RRID:AB_625558

Type: Antibody

Proper Citation

GeneTex Cat# GTX78382, RRID:AB_625558

Antibody Information

URL: http://antibodyregistry.org/AB_625558

Proper Citation: GeneTex Cat# GTX78382, RRID:AB_625558

Target Antigen: IgG Light chain [IVA285-1] antibody

Host Organism: mouse

Clonality: monoclonal

Comments: Discontinued; manufacturer recommendations: Flow Cytometry; Western Blot; Immunofluorescence; Immunohistochemistry; Immunohistochemistry - frozen; Flow cytometry/FACS, IHC (Frozen sections), Western blot. The usefulness of this product in other applications has not been determined., FACS, IHC-Fr, WB, WB: Recommended dilution:0,5-1 ?g/ml, overnight in 4°C Positive control and sample preparation:BS (bovine serum), dilution 1:50 in Laemmli reducing buffer, boiled in water bath for 3 min Application notes: Strongly reacts with bovine IgG light chains, weakly reacts with IgM. Reducing conditions. SDS-PAGE (12 separating gel, 4 stacking gel). Recommended secondary antibody - anti mouse IgG/HRP - dilution 1:2000 in PBS/Tween, 60 min on vertical incubator.

Antibody Name: IgG Light chain [IVA285-1] antibody

Description: This monoclonal targets IgG Light chain [IVA285-1] antibody

Target Organism: bovine - not yet tested in other species, bovine

Antibody ID: AB_625558

Vendor: GeneTex

Catalog Number: GTX78382

Record Creation Time: 20231110T080402+0000

Ratings and Alerts

No rating or validation information has been found for IgG Light chain [IVA285-1] antibody.

Warning: Discontinued at GeneTex

Discontinued; manufacturer recommendations: Flow Cytometry; Western Blot; Immunofluorescence; Immunohistochemistry; Immunohistochemistry - frozen; Flow cytometry/FACS, IHC (Frozen sections), Western blot. The usefulness of this product in other applications has not been determined., FACS, IHC-Fr, WB, WB: Recommended dilution:0,5-1 µg/ml, overnight in 4°C Positive control and sample preparation:BS (bovine serum), dilution 1:50 in Laemmli reducing buffer, boiled in water bath for 3 min Application notes: Strongly reacts with bovine IgG light chains, weakly reacts with IgM. Reducing conditions. SDS-PAGE (12 separating gel, 4 stacking gel). Recommended secondary antibody - anti mouse IgG/HRP - dilution 1:2000 in PBS/Tween, 60 min on vertical incubator.

Data and Source Information

Source: [Antibody Registry](#)

Usage and Citation Metrics

We have not found any literature mentions for this resource.