

Resource Summary Report

Generated by [RRID](#) on Sep 9, 2026

ANILLIN / Scraps (C Terminus) antibody

RRID:AB_10721683

Type: Antibody

Proper Citation

GeneTex Cat# GTX89791, RRID:AB_10721683

Antibody Information

URL: http://antibodyregistry.org/AB_10721683

Proper Citation: GeneTex Cat# GTX89791, RRID:AB_10721683

Target Antigen: ANILLIN / Scraps (C Terminus) antibody

Host Organism: goat

Clonality: polyclonal

Comments: Discontinued; manufacturer recommendations: ELISA; Immunohistochemistry; Preliminary experiments gave an approx 75kDa band in Mouse Brain lysates after 0.3µg/ml antibody staining. Please note that currently we cannot find an explanation in the literature for the band we observe given the calculated size of 124kDa according to NP_061155.2. The 75kDa band was successfully blocked by incubation with the immunizing peptide. We would appreciate any feedback from people in the field - have any results been reported with other antibodies/lysates? Have any further splice variants/modified forms been reported? In paraffin embedded Human Kidney shows staining of nuclei in some cells of renal tubules.. Recommended concentration, 3-10µg/ml. ELISA, Immunohistochemistry.

Antibody Name: ANILLIN / Scraps (C Terminus) antibody

Description: This polyclonal targets ANILLIN / Scraps (C Terminus) antibody

Target Organism: cow, mouse, bovine, human

Antibody ID: AB_10721683

Vendor: GeneTex

Catalog Number: GTX89791

Record Creation Time: 20231110T065849+0000

Ratings and Alerts

No rating or validation information has been found for ANILLIN / Scraps (C Terminus) antibody.

Warning: Discontinued at GeneTex

Discontinued; manufacturer recommendations: ELISA; Immunohistochemistry; Preliminary experiments gave an approx 75kDa band in Mouse Brain lysates after 0.3µg/ml antibody staining. Please note that currently we cannot find an explanation in the literature for the band we observe given the calculated size of 124kDa according to NP_061155.2. The 75kDa band was successfully blocked by incubation with the immunizing peptide. We would appreciate any feedback from people in the field - have any results been reported with other antibodies/lysates? Have any further splice variants/modified forms been reported? In paraffin embedded Human Kidney shows staining of nuclei in some cells of renal tubules.. Recommended concentration, 3-10µg/ml. ELISA, Immunohistochemistry.

Data and Source Information

Source: [Antibody Registry](#)

Usage and Citation Metrics

We have not found any literature mentions for this resource.