

Resource Summary Report

Generated by [RRID](#) on Aug 14, 2026

Ymer / CCDC50 antibody

RRID:AB_10723066

Type: Antibody

Proper Citation

GeneTex Cat# GTX89553, RRID:AB_10723066

Antibody Information

URL: http://antibodyregistry.org/AB_10723066

Proper Citation: GeneTex Cat# GTX89553, RRID:AB_10723066

Target Antigen: Ymer / CCDC50 antibody

Host Organism: goat

Clonality: polyclonal

Comments: Discontinued; manufacturer recommendations: Immunohistochemistry; ELISA; Preliminary experiments gave an approx 18kDa band in Human Brain lysates after 1µg/ml antibody staining. Please note that currently we cannot find an explanation in the literature for the band we observe given the calculated size of 56.3kDa according to NP_848018.1 and 35.8kDa according to NP_777568.1. The 18kDa band was successfully blocked by incubation with the immunizing peptide. We would appreciate any feedback from people in the field - have any results been reported with other antibodies/lysates? Have any further splice variants/modified forms been reported? In paraffin embedded Human Pancreas shows strong staining in cytoplasm in select acinar cells. Recommended concentration, 5-10µg/ml. This antibody is expected to recognise both reported

Antibody Name: Ymer / CCDC50 antibody

Description: This polyclonal targets Ymer / CCDC50 antibody

Target Organism: rat, canine, cow, bovine, dog, human

Antibody ID: AB_10723066

Vendor: GeneTex

Catalog Number: GTX89553

Record Creation Time: 20250820T020755+0000

Ratings and Alerts

No rating or validation information has been found for Ymer / CCDC50 antibody.

Warning: Discontinued at GeneTex

Discontinued; manufacturer recommendations: Immunohistochemistry; ELISA; Preliminary experiments gave an approx 18kDa band in Human Brain lysates after 1µg/ml antibody staining. Please note that currently we cannot find an explanation in the literature for the band we observe given the calculated size of 56.3kDa according to NP_848018.1 and 35.8kDa according to NP_777568.1. The 18kDa band was successfully blocked by incubation with the immunizing peptide. We would appreciate any feedback from people in the field - have any results been reported with other antibodies/lysates? Have any further splice variants/modified forms been reported? In paraffin embedded Human Pancreas shows strong staining in cytoplasm in select acinar cells. Recommended concentration, 5-10µg/ml. This antibody is expected to recognise both reported

Data and Source Information

Source: [Antibody Registry](#)

Usage and Citation Metrics

We have not found any literature mentions for this resource.