

Resource Summary Report

Generated by [RRID](#) on Jul 28, 2026

CD44 antibody [IM7] (FITC)

RRID:AB_11163977

Type: Antibody

Proper Citation

GeneTex Cat# GTX80087, RRID:AB_11163977

Antibody Information

URL: http://antibodyregistry.org/AB_11163977

Proper Citation: GeneTex Cat# GTX80087, RRID:AB_11163977

Target Antigen: CD44 antibody [IM7] (FITC)

Host Organism: rat

Clonality: monoclonal

Comments: Discontinued; manufacturer recommendations: Immunohistochemistry; Immunoprecipitation; Immunohistochemistry - fixed; Flow Cytometry; Immunocytochemistry; Immunofluorescence; Western Blot; Immunohistochemistry - frozen; Flow cytometry, Immunocytochemistry, Immunofluorescence, Immunohistochemistry, IHC (Frozen sections), IHC (Formalin-fixed paraffin-embedded sections), Immunoprecipitation, Western Blot. The usefulness of this product in other applications has not been determined., FACS, ICC, IF, IHC, IHC-Fr, IHC-P, IP, WB, Recommended Starting Dilutions: FACS: Use at 1 µg/mL. Positive Control:mouse peripheral blood; IHC-P: Positive tissue:mouse spleen; Optimal dilutions should be determined experimentally by the researcher.

Antibody Name: CD44 antibody [IM7] (FITC)

Description: This monoclonal targets CD44 antibody [IM7] (FITC)

Target Organism: feline, canine, horse, mouse, cat, dog, human

Antibody ID: AB_11163977

Vendor: GeneTex

Catalog Number: GTX80087

Record Creation Time: 20250820T042824+0000

Ratings and Alerts

No rating or validation information has been found for CD44 antibody [IM7] (FITC).

Warning: Discontinued at GeneTex

Discontinued; manufacturer recommendations: Immunohistochemistry; Immunoprecipitation; Immunohistochemistry - fixed; Flow Cytometry; Immunocytochemistry; Immunofluorescence; Western Blot; Immunohistochemistry - frozen; Flow cytometry, Immunocytochemistry, Immunofluorescence, Immunohistochemistry, IHC (Frozen sections), IHC (Formalin-fixed paraffin-embedded sections), Immunoprecipitation, Western Blot. The usefulness of this product in other applications has not been determined., FACS, ICC, IF, IHC, IHC-Fr, IHC-P, IP, WB, Recommended Starting Dilutions: FACS: Use at 1 µg/mL. Positive Control:mouse peripheral blood; IHC-P: Positive tissue:mouse spleen; Optimal dilutions should be determined experimentally by the researcher.

Data and Source Information

Source: [Antibody Registry](#)

Usage and Citation Metrics

We have not found any literature mentions for this resource.