

Resource Summary Report

Generated by [RRID](#) on Jul 29, 2026

CD38 antibody [HIT2] (PE)

RRID:AB_11167028

Type: Antibody

Proper Citation

GeneTex Cat# GTX80118, RRID:AB_11167028

Antibody Information

URL: http://antibodyregistry.org/AB_11167028

Proper Citation: GeneTex Cat# GTX80118, RRID:AB_11167028

Target Antigen: CD38 antibody [HIT2] (PE)

Host Organism: mouse

Clonality: monoclonal

Comments: Discontinued; manufacturer recommendations: Flow Cytometry; Immunofluorescence; Immunohistochemistry; Immunohistochemistry - frozen; Immunohistochemistry - fixed; Western Blot; Recommended Starting Dilutions: WB: Use at 2 µg/mL. Positive Control: RAJI human cell line Sample preparation: Resuspend approx. 50 mil. cells in 1 ml cold Lysis buffer (1 laurylmaltoside in 20 mM Tris/Cl, 100 mM NaCl pH 8.2, 50 mM NaF including Protease inhibitor Cocktail). Incubate 60 min on ice. Centrifuge to remove cell debris. Mix lysate with non-reducing SDS-PAGE Sample buffer. Boil for 5 min. Use under non-reducing conditions; IHC-P: Use at 10 µg/mL; Optimal dilutions should be determined experimentally by the researcher., Flow cytometry, Immunohistochemistry, IHC (Frozen sections), IHC (Formalin-fixed paraffin-embedded sections), Western

Antibody Name: CD38 antibody [HIT2] (PE)

Description: This monoclonal targets CD38 antibody [HIT2] (PE)

Target Organism: human

Antibody ID: AB_11167028

Vendor: GeneTex

Catalog Number: GTX80118

Record Creation Time: 20250819T215356+0000

Record Last Update: 20250820T002720+0000

Ratings and Alerts

No rating or validation information has been found for CD38 antibody [HIT2] (PE).

Warning: Discontinued at GeneTex

Discontinued; manufacturer recommendations: Flow Cytometry; Immunofluorescence; Immunohistochemistry; Immunohistochemistry - frozen; Immunohistochemistry - fixed; Western Blot; Recommended Starting Dilutions: WB: Use at 2 µg/mL. Positive Control: RAJI human cell line Sample preparation: Resuspend approx. 50 mil. cells in 1 ml cold Lysis buffer (1 laurylmaltoside in 20 mM Tris/Cl, 100 mM NaCl pH 8.2, 50 mM NaF including Protease inhibitor Cocktail). Incubate 60 min on ice. Centrifuge to remove cell debris. Mix lysate with non-reducing SDS-PAGE Sample buffer. Boil for 5 min. Use under non-reducing conditions; IHC-P: Use at 10 µg/mL; Optimal dilutions should be determined experimentally by the researcher., Flow cytometry, Immunohistochemistry, IHC (Frozen sections), IHC (Formalin-fixed paraffin-embedded sections), Western

Data and Source Information

Source: [Antibody Registry](#)

Usage and Citation Metrics

We have not found any literature mentions for this resource.