

# Resource Summary Report

Generated by [RRID](#) on Aug 30, 2026

## Vimentin antibody [VI-01] (Dyomics 547)

RRID:AB\_11174925

Type: Antibody

---

### Proper Citation

GeneTex Cat# GTX79853, RRID:AB\_11174925

---

### Antibody Information

**URL:** [http://antibodyregistry.org/AB\\_11174925](http://antibodyregistry.org/AB_11174925)

**Proper Citation:** GeneTex Cat# GTX79853, RRID:AB\_11174925

**Target Antigen:** Vimentin antibody [VI-01] (Dyomics 547)

**Host Organism:** mouse

**Clonality:** monoclonal

**Comments:** Discontinued; manufacturer recommendations: Recommended Starting Dilutions: ICC/IF: Staining technique:RBL rat basophilic leukemia cell line:(a) Fix cells for 10 min in methanol at -20 degree and for 6 min in acetone at -20 degree; (b) Fix cells directly in methanol for 10 min at -20 degree or in acetone for 10 min at -20 degree.Incubation: 45 min RT. Positive Control:3T3 mouse Swiss albino fibroblast cell lineRBL rat basophilic leukemia cell line; Optimal dilutions should be determined experimentally by the researcher., Immunocytochemistry, Immunofluorescence, Western Blot. The usefulness of this product in other applications has not been determined., ICC, IF, WB; Western Blot; Immunofluorescence; Immunocytochemistry

**Antibody Name:** Vimentin antibody [VI-01] (Dyomics 547)

**Description:** This monoclonal targets Vimentin antibody [VI-01] (Dyomics 547)

**Target Organism:** feline, rat, hamster, donkey, porcine, canine, goat, horse, mammalian, mouse, rabbit, bovine, human, sheep

**Antibody ID:** AB\_11174925

**Vendor:** GeneTex

**Catalog Number:** GTX79853

**Record Creation Time:** 20250819T230134+0000

## Ratings and Alerts

No rating or validation information has been found for Vimentin antibody [VI-01] (Dyomics 547).

**Warning:** Discontinued at GeneTex

Discontinued; manufacturer recommendations: Recommended Starting Dilutions: ICC/IF: Staining technique:RBL rat basophilic leukemia cell line:(a) Fix cells for 10 min in methanol at -20 degree and for 6 min in acetone at -20 degree; (b) Fix cells directly in methanol for 10 min at -20 degree or in acetone for 10 min at -20 degree.Incubation: 45 min RT. Positive Control:3T3 mouse Swiss albino fibroblast cell lineRBL rat basophilic leukemia cell line; Optimal dilutions should be determined experimentally by the researcher., Immunocytochemistry, Immunofluorescence, Western Blot. The usefulness of this product in other applications has not been determined., ICC, IF, WB; Western Blot; Immunofluorescence; Immunocytochemistry

---

## Data and Source Information

**Source:** [Antibody Registry](#)

---

## Usage and Citation Metrics

We have not found any literature mentions for this resource.