

Resource Summary Report

Generated by [RRID](#) on Aug 2, 2026

ESA [ESA214] antibody (FITC)

RRID:AB_1240769

Type: Antibody

Proper Citation

GeneTex Cat# GTX30708, RRID:AB_1240769

Antibody Information

URL: http://antibodyregistry.org/AB_1240769

Proper Citation: GeneTex Cat# GTX30708, RRID:AB_1240769

Target Antigen: ESA [ESA214] antibody (FITC)

Host Organism: mouse

Clonality: monoclonal

Comments: Discontinued; manufacturer recommendations: Immunohistochemistry; Flow Cytometry; Immunohistochemistry - frozen; Immunofluorescence; Immunocytochemistry; Immunohistochemistry - fixed; Flow cytometry/FACS, Immunocytochemistry, Immunofluorescence, IHC (Frozen sections), IHC (Formalin-fixed paraffin-embedded sections). The usefulness of this product in other applications has not been determined., FACS, ICC, IF, IHC-Fr, IHC-P, For FACS: Use at a dilution of 1:200-1:1,000 (Not verified by in-house testing). For ICC: Use at an assay dependent dilution. For IF: Use at a dilution of 1:10-1:20. For IHC-Fr: Use at a dilution of 1:50-1:100. For IHC-P: Use at a dilution of 1:50-1:100. Not tested in other applications. Optimal dilutions/concentrations should be determined by the end user.

Antibody Name: ESA [ESA214] antibody (FITC)

Description: This monoclonal targets ESA [ESA214] antibody (FITC)

Antibody ID: AB_1240769

Vendor: GeneTex

Catalog Number: GTX30708

Record Creation Time: 20250820T010419+0000

Record Last Update: 20250820T095714+0000

Ratings and Alerts

No rating or validation information has been found for ESA [ESA214] antibody (FITC).

Warning: Discontinued at GeneTex

Discontinued; manufacturer recommendations: Immunohistochemistry; Flow Cytometry; Immunohistochemistry - frozen; Immunofluorescence; Immunocytochemistry; Immunohistochemistry - fixed; Flow cytometry/FACS, Immunocytochemistry, Immunofluorescence, IHC (Frozen sections), IHC (Formalin-fixed paraffin-embedded sections). The usefulness of this product in other applications has not been determined., FACS, ICC, IF, IHC-Fr, IHC-P, For FACS: Use at a dilution of 1:200-1:1,000 (Not verified by in-house testing). For ICC: Use at an assay dependent dilution. For IF: Use at a dilution of 1:10-1:20. For IHC-Fr: Use at a dilution of 1:50-1:100. For IHC-P: Use at a dilution of 1:50-1:100. Not tested in other applications. Optimal dilutions/concentrations should be determined by the end user.

Data and Source Information

Source: [Antibody Registry](#)

Usage and Citation Metrics

We found 3 mentions in open access literature.

Listed below are recent publications. The full list is available at [RRID](#) .

Verhagen MP, et al. (2024) The SW480 cell line as a model of resident and migrating colon cancer stem cells. iScience, 27(9), 110658.

Xu T, et al. (2022) Alternative splicing downstream of EMT enhances phenotypic plasticity and malignant behavior in colon cancer. eLife, 11.

Sacchetti A, et al. (2021) Phenotypic plasticity underlies local invasion and distant metastasis in colon cancer. eLife, 10.