

Resource Summary Report

Generated by [RRID](#) on Sep 11, 2026

CD105 [SN6h] antibody

RRID:AB_366904

Type: Antibody

Proper Citation

GeneTex Cat# GTX73783, RRID:AB_366904

Antibody Information

URL: http://antibodyregistry.org/AB_366904

Proper Citation: GeneTex Cat# GTX73783, RRID:AB_366904

Target Antigen: CD105 [SN6h] antibody

Host Organism: mouse

Clonality: monoclonal

Comments: Discontinued; manufacturer recommendations: IHC, IHC-P, Immunohistochemistry, IHC (Formalin-fixed paraffin-embedded sections). The usefulness of this product in other applications has not been determined., This antibody may be diluted to a titer of 1:10 -1:20 in an ABC method. However, depending on the staining conditions employed, we suggest that final dilution should be determined by the user. We suggest an incubation period of 60 minutes at room temperature. However, depending upon the fixation conditions and the staining system employed, optimal incubation should be determined by the end user. Formalin fixed paraffin embedded tissue sections require predigestion with Protease XXXV.; Immunohistochemistry - fixed; Immunohistochemistry

Antibody Name: CD105 [SN6h] antibody

Description: This monoclonal targets CD105 [SN6h] antibody

Antibody ID: AB_366904

Vendor: GeneTex

Catalog Number: GTX73783

Record Creation Time: 20231110T081250+0000

Record Last Update: 20260829T070146+0000

Ratings and Alerts

No rating or validation information has been found for CD105 [SN6h] antibody.

Warning: Discontinued at GeneTex

Discontinued; manufacturer recommendations: IHC, IHC-P, Immunohistochemistry, IHC (Formalin-fixed paraffin-embedded sections). The usefulness of this product in other applications has not been determined., This antibody may be diluted to a titer of 1:10 -1:20 in an ABC method. However, depending on the staining conditions employed, we suggest that final dilution should be determined by the user. We suggest an incubation period of 60 minutes at room temperature. However, depending upon the fixation conditions and the staining system employed, optimal incubation should be determined by the end user. Formalin fixed paraffin embedded tissue sections require predigestion with Protease XXXV.; Immunohistochemistry - fixed; Immunohistochemistry

Data and Source Information

Source: [Antibody Registry](#)

Usage and Citation Metrics

We have not found any literature mentions for this resource.