

Resource Summary Report

Generated by [RRID](#) on Sep 11, 2026

P Glycoprotein [C494] antibody

RRID:AB_367205

Type: Antibody

Proper Citation

GeneTex Cat# GTX23365, RRID:AB_367205

Antibody Information

URL: http://antibodyregistry.org/AB_367205

Proper Citation: GeneTex Cat# GTX23365, RRID:AB_367205

Target Antigen: P Glycoprotein [C494] antibody

Host Organism: mouse

Clonality: monoclonal

Comments: Discontinued; manufacturer recommendations: Western Blot; Immunocytochemistry; Immunohistochemistry - fixed; Immunoprecipitation; Immunohistochemistry; For ICC: Use at an assay dependent dilution. IHC-P: Use at a dilution of 1:20 or greater, for Biotin-based detection systems. For optimal staining, the primary antibody should be incubated 20-60 minutes at room temperature. No pre-treatment required. Note: Unequivocal plasma membrane patterns of immunostaining represent true P Glycoprotein expression. Weak homogeneous, cytoplasmic, or granular patterns of reactivity may represent staining of the Pyruvate Carboxylase cross-reactive epitope rather than positive staining for P Glycoprotein. IP: Use at an assay dependent dilution. WB: Use at an assay dependent dilution. Predicted molecular weight: 141 kDa. Not tested in other applicatio

Antibody Name: P Glycoprotein [C494] antibody

Description: This monoclonal targets P Glycoprotein [C494] antibody

Antibody ID: AB_367205

Vendor: GeneTex

Catalog Number: GTX23365

Record Creation Time: 20231110T081303+0000

Record Last Update: 20260831T221024+0000

Ratings and Alerts

No rating or validation information has been found for P Glycoprotein [C494] antibody.

Warning: Discontinued at GeneTex

Discontinued; manufacturer recommendations: Western Blot; Immunocytochemistry; Immunohistochemistry - fixed; Immunoprecipitation; Immunohistochemistry; For ICC: Use at an assay dependent dilution. IHC-P: Use at a dilution of 1:20 or greater, for Biotin-based detection systems. For optimal staining, the primary antibody should be incubated 20-60 minutes at room temperature. No pre-treatment required. Note: Unequivocal plasma membrane patterns of immunostaining represent true P Glycoprotein expression. Weak homogeneous, cytoplasmic, or granular patterns of reactivity may represent staining of the Pyruvate Carboxylase cross-reactive epitope rather than positive staining for P Glycoprotein. IP: Use at an assay dependent dilution. WB: Use at an assay dependent dilution. Predicted molecular weight: 141 kDa. Not tested in other applicatio

Data and Source Information

Source: [Antibody Registry](#)

Usage and Citation Metrics

We have not found any literature mentions for this resource.