

Resource Summary Report

Generated by [RRID](#) on Aug 4, 2026

Spectrin antibody [SB-SP1]

RRID:AB_367595

Type: Antibody

Proper Citation

GeneTex Cat# GTX11184, RRID:AB_367595

Antibody Information

URL: http://antibodyregistry.org/AB_367595

Proper Citation: GeneTex Cat# GTX11184, RRID:AB_367595

Target Antigen: Spectrin antibody [SB-SP1]

Host Organism: mouse

Clonality: monoclonal

Comments: Discontinued; manufacturer recommendations: IgG1; IgG1 Immunofluorescence; Immunohistochemistry; Western Blot; IF: Use at an assay dependent dilution. By immunofluorescence the antibody will stain human spectrin in erythrocyte smears. WB: Use at a dilution of 1/500 (determined using an extract of human erythrocyte ghosts or purified human erythrocyte spectrin and immunoperoxidase labeling). Expected bands at 220 kDa (beta) and 240 kDa (alpha) chains of human erythrocyte spectrin. The antibody stains the beta subunit stronger than the alpha subunit at the recommended titer. In addition, a number of polypeptides in the region of 170, 150, 85, and 45 kDa will react with the antibody. Not tested in other applications. Optimal dilutions/concentrations should be determined by the end user., IF, WB, Immunofluorescence, Western blot. The usefulness of

Antibody Name: Spectrin antibody [SB-SP1]

Description: This monoclonal targets Spectrin antibody [SB-SP1]

Antibody ID: AB_367595

Vendor: GeneTex

Catalog Number: GTX11184

Record Creation Time: 20250820T032634+0000

Record Last Update: 20250820T161548+0000

Ratings and Alerts

No rating or validation information has been found for Spectrin antibody [SB-SP1].

Warning: Discontinued at GeneTex

Discontinued; manufacturer recommendations: IgG1; IgG1 Immunofluorescence; Immunohistochemistry; Western Blot; IF: Use at an assay dependent dilution. By immunofluorescence the antibody will stain human spectrin in erythrocyte smears. WB: Use at a dilution of 1/500 (determined using an extract of human erythrocyte ghosts or purified human erythrocyte spectrin and immunoperoxidase labeling). Expected bands at 220 kDa (beta) and 240 kDa (alpha) chains of human erythrocyte spectrin. The antibody stains the beta subunit stronger than the alpha subunit at the recommended titer. In addition, a number of polypeptides in the region of 170, 150, 85, and 45 kDa will react with the antibody. Not tested in other applications. Optimal dilutions/concentrations should be determined by the end user., IF, WB, Immunofluorescence, Western blot. The usefulness of

Data and Source Information

Source: [Antibody Registry](#)

Usage and Citation Metrics

We have not found any literature mentions for this resource.