

Resource Summary Report

Generated by [RRID](#) on Aug 11, 2026

DC-SIGN antibody [120507]

RRID:AB_369176

Type: Antibody

Proper Citation

GeneTex Cat# GTX13487, RRID:AB_369176

Antibody Information

URL: http://antibodyregistry.org/AB_369176

Proper Citation: GeneTex Cat# GTX13487, RRID:AB_369176

Target Antigen: DC-SIGN antibody [120507]

Host Organism: mouse

Clonality: monoclonal

Comments: Discontinued; manufacturer recommendations: IgG2; IgG2 Flow cytometry/FACS, Inhibition assay. The usefulness of this product in other applications has not been determined., FACS, Inhib, FACS: Use at a concentration of 25 - 50 µg/ml (10 µl of a 25-50 µg/ml stock solution of the monoclonal antibody is mixed with up to 1 million cells in a minimal volume (0.2 ml) of buffer (PBS + 0.5 BSA). The reaction is incubated at room temperature for 30 minutes. Inhib: Use at a concentration of 5 µg/ml. (This antibody shows inhibition of DC-SIGN expressing NIH-3T3 cell adhesion to ICAM-3 coated 96 well plates. The adhesion of NIH-3T3-DC-SIGN cells (5 x 10⁴ cells/well) to plates coated with 5 µg/ml (100 µl/well) of recombinant human ICAM-3/Fc is maximally inhibited (100) using 100 µl of a 5 µg/ml antibody so

Antibody Name: DC-SIGN antibody [120507]

Description: This monoclonal targets DC-SIGN antibody [120507]

Antibody ID: AB_369176

Vendor: GeneTex

Catalog Number: GTX13487

Record Creation Time: 20250819T215043+0000

Record Last Update: 20250820T001714+0000

Ratings and Alerts

No rating or validation information has been found for DC-SIGN antibody [120507].

Warning: Discontinued at GeneTex

Discontinued; manufacturer recommendations: IgG2; IgG2 Flow cytometry/FACS, Inhibition assay. The usefulness of this product in other applications has not been determined., FACS, Inhib, FACS: Use at a concentration of 25 - 50 µg/ml (10 µl of a 25-50 µg/ml stock solution of the monoclonal antibody is mixed with up to 1 million cells in a minimal volume (0.2 ml) of buffer (PBS + 0.5 BSA). The reaction is incubated at room temperature for 30 minutes. Inhib: Use at a concentration of 5 µg/ml. (This antibody shows inhibition of DC-SIGN expressing NIH-3T3 cell adhesion to ICAM-3 coated 96 well plates. The adhesion of NIH-3T3-DC-SIGN cells (5 x 10⁴ cells/well) to plates coated with 5 µg/ml (100 µl/well) of recombinant human ICAM-3/Fc is maximally inhibited (100 %) using 100 µl of a 5 µg/ml antibody so

Data and Source Information

Source: [Antibody Registry](#)

Usage and Citation Metrics

We have not found any literature mentions for this resource.